
















































TABLE 1
SUMMARY OF OBSERVATIONS

Source (1)	z (2)	VLBI (3)	JCMT (4)	UKIRT (5)	ROSAT (6)	EGRET (7)
0016+731	1.76	Jun 1992 Sep 1992	Aug 1991 Oct 1992		Jul 1992 ...	Jul/Aug 1992 ...
0133+476	0.86	Jun 1992 Sep 1992	Aug 1991 Oct 1992	Aug 1992 ...	Aug 1992 ...	Nov/Dec 1991 ...
0212+735	2.37	Nov 1992 Mar 1993	Aug 1991 ...		Feb 1991 ...	Jul/Aug 1992 ...
NRAO 140	1.26	Jun 1991 Nov 1991	Aug 1991 Oct 1992	Aug 1992 ...	Aug 1992 ...	May 1991 ...
3C 111	0.05	Jun 1991 Nov 1991	Aug 1991 Oct 1992		Mar 1991 Feb 1993	Nov/Dec 1991 ...
0429+415	0.41	Jun 1992 Sep 1992	Aug 1991 Oct 1992			
3C 120	0.03	Jun 1991 Nov 1991	Aug 1991 Oct 1992		Feb 1991 ...	Aug 1992 ...
0552+398	2.36	Jun 1992 Sep 1992	Aug 1991 Oct 1992		Mar 1993 ...	June 1992 ...
0642+449	3.40	Jun 1991 ...	Aug 1991 Jan 1992		Mar 1991 ...	
0736+017	0.19	Jun 1991 ...	Aug 1991 Oct 1992		Apr 1991 Apr 1992	Nov 1992 ...
0804+499	1.43	Jun 1992 Sep 1992	Aug 1991 Oct 1992	Nov 1993 Mar 1994	Oct 1992 ...	Jan 1992 ...
0820+560	1.42	...	Aug 1991	Dec 1993	Oct 1993	...
0836+710	2.17	Jun 1992 Sep 1992	Mar 1992 Oct 1992		Mar 1992 Nov 1992	Jan 1992 Mar 1992
0917+449	2.18	Jan 1995	...	...	Oct/Nov 1992	...
4C 39.25	0.70		Aug 1991 Oct 1992	Nov 1993 ...	Apr 1991 Apr 1993	Sep/Oct 1992 ...
0945+408	1.25	Nov 1992 Mar 1993 Jan 1995	Oct 1992	May 1994 May 1994 ...	Nov 1993 Nov 1993 ...	Sept/Oct 1992 Sep/Oct 1993 ...
0954+556	0.91	...	Oct 1992	Dec 1993	Nov 1993	Jun/Jul 1991
0955+476	1.87	Jan 1995	Oct 1992	Dec 1993	Nov 1993	...
1150+497	0.334	 Jan 1995	Mar 1992 Oct 1992 ...	Jun 1992		Jun/Jul 1992
1150+812	1.25		Mar 1992		Apr 1992 May 1992 Oct 1992	Jan 1991
1611+343	1.40	Nov 1992 ... May 1996	Oct 1992	Aug 1992	Sep 1992 Jan 1993 ...	Sep 1992
1633+382	1.81	Sep 1992 Mar 1993	Oct 1992 ...	Aug 1992 ...	Aug 1992 ...	Nov 1992 Sep 1993
1638+398	1.66	...	Oct 1992	Sep 1992	Sep 1991	...
1641+399	0.60	Nov 1992 Mar 1993	Oct 1992 ...		Jun 1990 Jan 1992	Sep 1991 ...


TABLE 1—*Continued*

Source (1)	z (2)	VLBI (3)	JCMT (4)	UKIRT (5)	ROSAT (6)	EGRET (7)
1828+487	0.69	...	Aug 1991	...	...	Sep 1991
1954+513	1.22	Jan 1995	Aug 1991	...	...	May/Jun 1991
2005+403	1.74	Jan 1995	Aug 1991	...	...	May/Jun 1991
2037+511	1.69	...	Aug 1991	...	...	May/Jun 1991
		...	Oct 1992	...	...	...
		Jan 1995	...	...	...	...
2136+141	2.43	Jun 1991	Aug 1991	...	May 1991	Jan/Feb 1992
		Nov 1991	...	...	...	...
2201+315	0.30	Jan 1995	Aug 1991	...	Dec 1993	May/Jun 1991


Clean map. Array: BFHKLMNOPS
0820+560 at 22.225 GHz 1995 Jan 04




Clean map. Array: BFHKLMNOPS
0917+449 at 22.225 GHz 1995 Jan 04




Clean map. Array: BFHKLMNOPS
0945+408 at 22.225 GHz 1995 Jan 04




Clean map. Array: BFHKLMNOPS
0955+476 at 22.225 GHz 1995 Jan 04




Clean map. Array: BFHKLMNOPS
1150+497 at 22.225 GHz 1995 Jan 04




Clean map. Array: BFHKLMNOPS
1611+343 at 22.217 GHz 1996 May 04




Clean map. Array: BFHKLMNOPS
1954+513 at 22.225 GHz 1995 Jan 04



Clean map. Array: BFHKLMNOPS
2005+403 at 22.225 GHz 1995 Jan 04



Clean map. Array: BFHKLMNOPS
2037+511 at 22.225 GHz 1995 Jan 04



Clean map. Array: BFHKLMNPS
2201+315 at 22.225 GHz 1995 Jan 04




TABLE 2
VLBI ANTENNAS

Observatory	Location	Diameter (m)	Sessions ^a
Onsala Space Observatory	Onsala, Sweden	20	1,2,3,4,5,6
Max Plank-Institut für Radioastronomie	Effelsberg, Germany	100	1,2,3,4,5,6
Instituto di Radioastronomia	Medicina, Italy	32	1,2,3,4,5,6
Instituto di Radioastronomia	Noto, Italy	32	1,2,3,4,5,6
Metsähovi Radio Research Station	Metsähovi, Finland	14	3,4
Haystack Observatory	Westford, MA	37	1,2,3
Owens Valley Radio Observatory	Big Pine, CA	40	1,2
National Radio Astronomy Observatory	Green Bank, WV	43	1,2,3,4,5,6
National Radio Astronomy Observatory VLBA	Hancock, NH	25	5,6,7,8,9
National Radio Astronomy Observatory VLBA	Saint Croix	25	6,7,8,9
National Radio Astronomy Observatory VLA	Socorro, NM	25	1,2,3,4,5,6
National Radio Astronomy Observatory VLBA	Pie Town, NM	25	1,2,3,4,5,6,7,8,9
National Radio Astronomy Observatory VLBA	Los Alamos, NM	25	1,2,3,4,5,6,7,8,9
National Radio Astronomy Observatory VLBA	Fort Davis, TX	25	1,2,3,7,8,9
National Radio Astronomy Observatory VLBA	Kitt Peak, AZ	25	1,2,3,4,5,6,7,8,9
National Radio Astronomy Observatory VLBA	North Liberty, IA	25	1,2,3,4,5,6,7,8,9
National Radio Astronomy Observatory VLBA	Owens Valley, CA	25	3,4,5,6,7,8,9
National Radio Astronomy Observatory VLBA	Brewster, WA	25	3,4,5,6,7,8,9
National Radio Astronomy Observatory VLBA	Mauna Kea, HI	25	7,8,9

^aSession 1:1991.45, Session 2:1991.85, Session 3: 1992.45, Session 4: 1992.70, Session 5:1992.85, Session 6: 1993.18, Session 7:1995.01, Session 8:1995.34, Session 9:1996.34




TABLE 3
VLBI MAP PARAMETERS

Source	Frequency (GHz)	a_{beam} (mas)	b_{beam} (mas)	θ_{beam} (degrees)	Peak (Jy beam ⁻¹)	Contour Levels (% of Peak)
0016+731	8.4	0.54	0.39	5	0.93	1, 2, 3, 5, 10, 25, 75, 90
	22	0.50	0.15	5	0.56	5, 10, 25, 75, 90
0133+476	8.4	1.21	0.38	15	0.52	2, 3, 5, 10, 25, 75, 90
	22	0.70	0.15	7	1.5	5, 10, 25, 50, 75, 90
0212+735	8.4	0.83	0.38	20	1.2	2, 3, 5, 10, 25, 50, 75, 90
	22	0.25	0.18	75	0.50	10, 25, 50, 75, 90
NRAO140	8.4	3.04	0.46	5	1.0	5, 10, 25, 50, 75, 90
	22	0.88	0.18	-5	0.89	5, 10, 25, 50, 75, 90
3C 111	8.4	1.33	0.46	8	0.48	2, 3, 5, 10, 25, 50, 75, 90
	22	1.03	0.20	2	1.26	5, 10, 25, 50, 75, 90
0429+415	8.4	1.54	0.39	5	0.40	5, 10, 25, 50, 75, 90
3C 120	8.4	4.62	0.46	-10	0.43	5, 10, 25, 50, 75, 90
	22	1.43	0.20	-9	0.41	10, 25, 50, 75, 90
0552+398	8.4	1.39	0.39	-5	4.1	2, 3, 5, 10, 25, 50, 75, 90
	22	0.58	0.18	-5	2.5	3, 5, 10, 25, 50, 75, 90
0642+449	22	0.50	0.20	-14	1.7	3, 5, 10, 25, 50, 75, 90
0736+017	22	1.10	0.20	-10	1.4	10, 25, 50, 75, 90
0804+499	8.4	1.59	0.38	18	0.82	2, 3, 5, 10, 25, 50, 75, 90
	22	0.63	0.15	5	1.36	3, 5, 10, 25, 50, 75, 90
0820+560	22	0.446	0.308	-11.6	0.35	2, 4, 8, 16, 32, 64
0836+710	8.4	0.65	0.45	30	0.95	5, 10, 25, 50, 75, 90
	22	0.33	0.23	-60	1.6	3, 5, 10, 25, 50, 75, 90
0917+449	22	0.533	0.2750	-11.2	1.19	0.5, 1, 2, 4, 8, 16, 32, 64
0945+408	8.4	0.98	0.38	-7	1.49	2, 3, 5, 10, 25, 50, 75, 90
	22	0.49	0.12	-10	0.69	5, 10, 25, 50, 75, 90
	22	0.761	0.332	-27	0.60	2, 4, 8, 16, 32, 64
0955+476	22	0.539	0.273	-12.1	0.93	1, 2, 4, 8, 16, 32, 64
1150+497	22	0.528	0.263	-18.4	1.50	-0.25, 0.25, 0.5, 1, 2, 4, 8, 16, 32, 64
1611+343	22	0.48	0.15	-5	1.28	3, 5, 10, 25, 50, 75, 90
	22	0.696	0.328	-4.4	2.03	4, 8, 16, 32, 64, 90
1633+382	8.4	1.13	0.53	-7	1.153	2, 3, 5, 10, 25, 50, 75, 90
	22	0.61	0.12	-20	0.53	5, 10, 25, 50, 75, 90
3C 345	8.4	1.1	0.53	-10	6.768	3, 5, 10, 25, 50, 75, 90
	22	0.75	0.15	-10	4.5	5, 10, 25, 50, 75, 90
1954+513	22	0.629	0.265	12.7	0.69	1, 2, 4, 8, 16, 32, 64
2005+403	22	0.921	0.862	49.9	1.35	1, 2, 4, 8, 16, 32, 64
2037+511	22	0.588	0.266	3.7	1.80	0.5, 1, 2, 4, 8, 16, 32, 64
2136+141	8.4	4.86	0.53	-10	1.75	2, 3, 5, 10, 25, 50, 75, 90
	22	1.5	0.25	-10	1.47	3, 5, 10, 25, 50, 75, 90
2201+315	22	0.626	0.255	-17.4	1.00	0.5, 1, 2, 4, 8, 16, 32, 64




TABLE 4
VLBI GAUSSIAN MODEL PARAMETERS

Source	Frequency (GHz)	Flux Density (Jy)	r (mas)	θ_{pos} (degrees)	a (mas)	b/a	ϕ (degrees)	A. F. (amplitude)	A. F. (closure phase)
0016+731	8.4	1.61 ± 0.05	0.0	0.0	0.49	0.58	-54.0	3.37	1.03
	22	1.23 ± 0.06	0.0	0.0	0.29	0.60	8.1	1.24	1.21
0133+476	8.4	0.56 ± 0.06	0.0	0.0	0.24	0.50	-4.7	3.13	1.11
		0.14 ± 0.02	1.98	-55.2	1.68	0.56	119.0		
	22	1.66 ± 0.07	0.0	0.0	0.12	0.14	-0.3	1.28	1.09
0212+735	8.4	1.70 ± 0.02	0.0	0.0	0.42	0.52	-42.1	2.08	2.07
		0.38 ± 0.01	2.22	102.6	2.58	0.82	-92.6		
	22	1.54 ± 0.06	0.0	0.0	0.32	0.40	-44	1.20	1.31
NRAO140	8.4	1.16 ± 0.04	0.0	0.0	0.28	0.93	20.6	3.80	1.81
		0.61 ± 0.02	2.03	120	3.64	0.61	5.5		
	22	1.53 ± 0.21	0.0	0.0	0.36	0.65	-8.4	1.98	1.1
		0.19 ± 0.03	1.60	142	2.89	0.53	95		
3C 111	8.4	0.66 ± 0.03	0.0	0.0	0.93	0.38	14.7	1.60	1.19
		0.31 ± 0.01	0.84	66.0	3.48	0.36	61.23		
	22	1.45 ± 0.23	0.0	0.0	0.33	0.72	16.5	1.4	1.2
		0.82 ± 0.12	1.1	61	2.69	0.81	121		
0429+415	8.4	1.05 ± 0.04	0.0	0.0	1.67	0.37	1.8	5.03	2.9
3C 120	8.4	0.58 ± 0.04	0.0	0.0	2.1	0.16	-11.1	2.65	1.82
		1.21 ± 0.05	3.16	255.1	3.9	0.73	37.3		
		0.27 ± 0.01	7.03	249.8	4.07	0.38	-15.1		
	22	0.61 ± 0.21	0.0	0.0	0.35	0.53	12	1.27	1.15
		0.47 ± 0.16	2.5	-109	0.88	1.0	69		
		0.34 ± 0.11	5.5	-108	1.4	1.0	88		
0552+398	8.4	6.95 ± 0.23	0.0	0.0	0.60	0.82	-7.3	3.17	1.59
	22	5.88 ± 0.34	0.0	0.0	0.30	0.77	-6.0	2.05	1.12
0642+449	22	2.55 ± 0.27	0.0	0.0	0.21	0.71	-91.6	1.06	1.27
0736+017	22	1.58 ± 0.47	0.0	0.0	0.07	0.36	29	0.87	1.27
		0.45 ± 0.14	0.52	-78	0.41	0.48	-19		
		0.16 ± 0.05	2.30	-83	3.15	0.001	48		
0804+499	8.4	0.97 ± 0.06	0.0	0.0	0.40	0.60	7.04	1.60	1.11
		0.09 ± 0.05	1.82	72.0	4.02	0.13	11.0		
	22	1.67 ± 0.25	0.0	0.0	0.27	0.19	16.8	1.00	1.40
0820+560	22	0.44 ± 0.07	0.0	0.00	0.25	0.23	-67.5		
		0.21 ± 0.03	0.78	105.6	2.97	0.13	-42.1		
0836+710	8.4	1.44 ± 0.01	0.0	0.0	0.35	0.46	61.6	2.76	2.28
		0.47 ± 0.01	2.33	210.0	1.65	0.61	14.5		
	22	1.77 ± 0.13	0.0	0.0	0.18	0.26	32.1	1.64	1.07
0917+449	22	1.31 ± 0.07	0.00	0.0	0.20	0.50	7.4	0.27	0.21
		0.23 ± 0.01	1.13	174.4	0.99	0.48	14.6		
0945+408	8.4	1.75 ± 0.25	0.0	0.0	0.27	0.38	-74.6	2.43	1.43
	22	1.57 ± 0.03	0.0	0.0	0.34	0.59	-20.5	1.05	1.07
	22	1.09 ± 0.05	0.13	105.0	0.64	0.25	-82.8	0.358	0.304
0955+476	22	1.01 ± 0.05	0.00	0.0	0.12	0.89	79.17	0.742	0.266
1150+497	22	1.72 ± 0.09	0.00	0.0	0.19	0.22	19.9	1.132	0.101
		0.18 ± 0.01	0.61	-155.5	1.32	0.50	31.2		
1611+343	22	1.56 ± 0.34	0.0	0.0	0.24	0.38	1.4	4.4	1.33
		0.26 ± 0.06	0.45	1.68	0.33	0.001	5.3		

TABLE 4—*Continued*

Source	Frequency (GHz)	Flux Density (Jy)	r (mas)	θ_{pos} (degrees)	a (mas)	b/a	ϕ (degrees)	A. F. (amplitude)	A. F. (closure phase)
1633+382	22	1.86 ± 0.09	0.08	106.7	0.22	0.37	−5.7	0.309	0.186
		0.57 ± 0.03	2.81	171.1	1.61	0.49	−38.7		
		0.57 ± 0.03	0.15	37.1	6.86	0.17	50.0		
		0.09 ± 0.01	0.86	93.8	1.19	0.10	−21.8		
	8.4	1.60 ± 0.04	0.0	0.0	0.52	0.0	84.6	1.29	1.20
		0.41 ± 0.01	0.84	−75.7	0.95	0.89	105.7		
3C 345	22	0.79 ± 0.11	0.0	0.0	0.24	0.62	37	1.52	1.29
		1.18 ± 0.18	2.6	−32	3.16	0.40	−32		
	8.4	9.41 ± 0.09	0.0	0.0	0.50	0.12	87.5	3.56	3.65
		9.26 ± 0.28	0.0	0.0	0.33	0.78	2.9		
		2.39 ± 0.07	0.29	−87	0.17	0.59	−6.4		
1954+513	22	0.72 ± 0.11	0.00	0.0	0.103	0.10	66.3	0.438	0.190
		0.41 ± 0.06	0.48	−53.5	0.36	0.57	−21.4		
2005+403	22	1.59 ± 0.08	0.0	0.0	0.80	0.14	83.0	0.918	0.498
		0.35 ± 0.02	0.51	75.4	0.83	0.20	60.2		
2037+511	22	1.98 ± 0.10	0.00	0.00	0.18	0.52	−6.5	0.55	0.331
		0.68 ± 0.03	0.75	−157.1	0.88	0.37	83.8		
2136+141	8.4	1.70 ± 0.03	0.0	0.0	0.14	0.65	−60.5	4.63	1.23
		0.45 ± 0.01	0.52	−133	1.96	0.94	−24.4		
	22	1.89 ± 0.21	0.0	0.0	0.33	0.5	−23	1.4	1.0
2201+315	22	0.49 ± 0.05	0.85	−49	5.7	0.4	−10	0.852	0.457
		1.26 ± 0.06	0.0	0.0	0.23	0.25	31.3		
		0.48 ± 0.02	1.87	−142.9	2.20	0.28	42.2		





TABLE 5
MILLIMETER/SUBMILLIMETER OBSERVATIONS WITH THE JCMT ^c

Source	Date	Wavelength (μm)	Flux Density (Jy)	Variable? ^d
(1)	(2)	(3)	(4)	(5)
0736+017	1993 Nov 21	800	1.13 $\pm$ 0.08	0.2
		1100	1.27 $\pm$ 0.08	0.3
		1300	1.48 $\pm$ 0.11	...
		2000	1.70 $\pm$ 0.10	0.3
0917+449	1993 Dec 08	800	0.84 $\pm$ 0.10	No
		1100	0.95 $\pm$ 0.10	No
		1300	1.01 $\pm$ 0.10	...
		2000	1.19 $\pm$ 0.15	No
0923+392 (4C39.25)	1993 Dec 08	800	0.48 $\pm$ 0.06	No
		1100	2.46 $\pm$ 0.17	No
		1300	2.79 $\pm$ 0.15	...
		2000	3.62 $\pm$ 0.45	No
0945+408	1993 Dec 08	800	<0.39	No
		1100	...	No
		1300	0.39 $\pm$ 0.10	No
		2000	0.59 $\pm$ 0.13	No
0954+556	1993 Dec 08	800	<0.23	No
		1100	0.16 $\pm$ 0.04	No
		1300	0.20 $\pm$ 0.04	No
		2000	0.47 $\pm$ 0.08	No
0955+476	1993 Dec 08	800	0.21 $\pm$ 0.07	No
		1100	0.23 $\pm$ 0.05	No
		1300	0.40 $\pm$ 0.05	No
		2000	0.44 $\pm$ 0.08	No
1641+399 (3C 345)	1993 Sep 14	800	1.35 $\pm$ 0.12	0.4
		1100	1.85 $\pm$ 0.07	0.3
		1300	2.24 $\pm$ 0.10	...
		2000	2.91 $\pm$ 0.15	0.3
2005+403	1993 Sep 14	800	0.23 $\pm$ 0.07	...
		1100	0.29 $\pm$ 0.05	0.3
		1300	0.47 $\pm$ 0.08	No
		2000	0.70 $\pm$ 0.14	No
2201+315	1993 Sep 14	1100	0.40 $\pm$ 0.07	0.3
		1300	0.41 $\pm$ 0.06	0.3
		2000	0.60 $\pm$ 0.13	0.3

^csee Bloom *et al.* 1994 for data from earlier epochs

^d $\frac{F_{max}-F_{min}}{F_{max}+F_{min}}$; “No”, if variability is not significant, as defined in text




TABLE 6
INFRARED OBSERVATIONS WITH UKIRT ^a

Source	Date	Wavelength (μm)	Flux Density (mJy)	α_{IR}	Instrument
(1)	(2)	(3)	(4)	(5)	(6)
0804+499	1993 Nov 19	1.25	1.70 $\pm$ 0.06	-0.01	UKT9
		1.65	1.81 $\pm$ 0.07		
		2.20	1.69 $\pm$ 0.04		
	1994 Mar 5	1.25	0.33 $\pm$ 0.01	0.06	UKT9
		1.65	0.41 $\pm$ 0.03		
		2.20	0.37 $\pm$ 0.03		
0820+560	1993 Dec 09	1.25	0.30 $\pm$ 0.02	0.63	UKT9
		1.65	0.42 $\pm$ 0.04		
		2.20	0.43 $\pm$ 0.03		
0923+392 (4C 39.25)	1993 Nov 19	1.25	1.59 $\pm$ 0.03	0.38	UKT9
		1.65	1.43 $\pm$ 0.03		
		2.20	1.97 $\pm$ 0.07		
0945+408	1994 May 13	1.25	0.33 $\pm$ 0.06	0.90	IRCAM
		1.65	0.45 $\pm$ 0.09		
		2.20	0.55 $\pm$ 0.11		
	1994 May 17	1.25	0.30 $\pm$ 0.06	0.97	IRCAM
		1.65	0.39 $\pm$ 0.08		
		2.20	0.52 $\pm$ 0.10		
0954+556	1993 Dec 09	1.25	0.77 $\pm$ 0.02	0.94	UKT9
		1.65	0.96 $\pm$ 0.03		
		2.20	1.31 $\pm$ 0.04		
0955+476	1993 Dec 09	1.25	0.71 $\pm$ 0.03	1.61	UKT9
		1.65	1.05 $\pm$ 0.05		
		2.20	1.76 $\pm$ 0.07		
1641+399 (3C 345)	1993 Sep 15	1.25	1.03 $\pm$ 0.01	1.51	UKT9
		1.65	1.48 $\pm$ 0.06		
		2.20	2.42 $\pm$ 0.06		
2005+403	1993 Sep 15	1.25	0.29 $\pm$ 0.02	0.66	UKT9
		1.65	0.33 $\pm$ 0.02		
		2.20	0.42 $\pm$ 0.04		
2201+315	1993 Sep 15	1.25	3.99 $\pm$ 0.03	1.52	UKT9
		1.65	5.48 $\pm$ 0.03		
		2.20	9.39 $\pm$ 0.05		

^asee Bloom *et al.* 1994 for data from earlier epochs




TABLE 7
X-RAY SPECTRAL PARAMETERS

Source	Date	Counts (0.1–2.5 keV)	Obs Time (sec)	$\log N_{\text{H,gal}}$ (atoms cm^{-2})	$\log N_{\text{H,best}}$ (atoms cm^{-2})	α	$F_{1\text{keV}}$ (μJy)	χ^2
0016+731	1992 Jul 29	40 ± 7	5243	21.38	...	1.0	0.06 ± 0.01	6.9
0133+476	1992 Aug 12	285 ± 17	4036	21.37	$20.92^{+0.08}_{-0.12}$	$0.66^{+0.64}_{-0.36}$	$0.27^{+0.10}_{-0.04}$	14
0212+735 ^a	1991 Feb 16	308 ± 18	7210	21.39	...	-0.60 ± 0.40	$0.20^{+0.02}_{-0.03}$	9
NRAO140 ^b	1992 Aug 8	921 ± 31	4039	21.41	21.46 ± 0.02	$1.01^{+0.16}_{-0.17}$	$1.59^{+1.11}_{-0.61}$	33
3C 111 ^b	1991 Mar 8	265 ± 17	838	22.08	...	$2.25^{+0.65}_{-0.40}$	$12.3^{+1.9}_{-2.2}$	7
	1993 Feb 13	840 ± 29	2322	22.08	21.98 ± 0.02	1.48 ± 0.20	$8.4^{+10.3}_{-4.4}$	16
3C 120	1991 Feb 20	3642 ± 62	1724	21.03	21.54 ± 0.02	$2.13^{+0.09}_{-0.11}$	$18.4^{+4.6}_{-3.9}$	36
0552+398	1993 Mar 13	185 ± 14	4294	21.28	...	$0.09^{+0.36}_{-0.16}$	0.21 ± 0.03	12
0642+449	1991 Mar 13	73 ± 10	3108	20.79	...	$0.79^{+0.86}_{-0.59}$	$0.08^{+0.01}_{-0.02}$	20
0736+017	1991 Apr 3	234 ± 16	1936	21.08	...	$1.24^{+0.36}_{-0.14}$	$0.50^{+0.06}_{-0.05}$	21
	1992 Apr 6	396 ± 20	3296	21.08	...	$1.45^{+0.25}_{-0.45}$	$0.47^{+0.06}_{-0.05}$	24
0804+499	1992 Oct 27	144 ± 12	2999	21.32	...	$1.09^{+0.71}_{-0.79}$	$0.24^{+0.11}_{-0.07}$	44
0836+710 ^a	1992 Mar 3	5365 ± 75	6990	20.46	20.54 ± 0.05	0.57 ± 0.11	2.37 ± 0.10	41
	1992 Nov 2	1904 ± 44	5030	20.46	20.56 ± 0.10	$0.57^{+0.19}_{-0.18}$	$1.21^{+0.08}_{-0.09}$	15
0917+449	1993 Oct 27-Nov 3	639 ± 28	3225	20.55	20.53 ± 0.10	$0.94^{+0.30}_{-0.24}$	$0.49^{+0.25}_{-0.20}$	40

TABLE 7—*Continued*

Source	Date	Counts (0.1–2.5 keV)	Obs Time (sec)	$\log N_{\mathrm{H,gal}}$ (atoms cm^{-2})	$\log N_{\mathrm{H,best}}$ (atoms cm^{-2})	α	$F_{1\mathrm{keV}}$ ($\mu\mathrm{Jy}$)	χ^2	D.
0923+392 ^c	1991 Apr 15	741	...	20.23	20.20 ± 0.20	$1.17^{+0.38}_{-0.34}$	0.52 ± 0.06	...	...
	1993 Apr 22	1356	...	20.23	$20.30^{+0.16}_{-0.20}$	1.32 ± 0.30	$0.69^{+0.07}_{-0.06}$	...	...
0945+408	1993 Nov 10	189 ± 16	2577	20.16	...	$1.06^{+0.22}_{-0.24}$	0.09 ± 0.02	46	31
0954+556	1993 Nov 8	393 ± 24	4293	19.95	$19.92^{+0.28}_{-0.40}$	$1.30^{+0.45}_{-0.30}$	0.08 ± 0.02	53	31
0955+476	1993 Nov 13	339 ± 20	3450	20.01	...	$0.19^{+0.15}_{-0.26}$	$0.21^{+0.05}_{-0.04}$	45	31
1150+812 ^a	1992 Apr 7	58 ± 9	2930	20.65	...	$0.55^{+0.21}_{-0.25}$	$0.09^{+0.09}_{-0.10}$	13	5
	1992 May 7	172 ± 14	5650	20.65	...	(Combined)	...	...	...
	1992 Oct 29	241 ± 17	9500	20.65	...	$0.41^{+0.19}_{-0.29}$	$0.09^{+0.10}_{-0.09}$	5.4	4
1611+343	1992 Sep 30	418 ± 21	3651	20.45	$20.32^{+0.08}_{-0.12}$	1.34 ± 0.30	$0.15^{+0.03}_{-0.01}$	80	30
	1993 Jan 17	332 ± 18	3153	20.45	$20.30^{+0.15}_{-0.25}$	0.65 ± 0.35	$0.24^{+0.04}_{-0.03}$	16	24
1633+382	1992 Aug 20	1208 ± 38	5781	20.58	$20.18^{+0.12}_{-0.16}$	$0.76^{+0.17}_{-0.20}$	$0.41^{+0.04}_{-0.03}$	41	30
1638+398	1992 Sep 3	64 ± 9	2857	20.49	...	1.89 ± 0.30	0.04 ± 0.01	15	15
3C 345	1990 Jun 19	1193 ± 43	4086	20.56	$19.71^{+0.20}_{-0.40}$	$0.77^{+0.13}_{-0.17}$	$0.39^{+0.04}_{-0.03}$	54	30
	1992 Jun 16	1014 ± 40	3899	20.56	$20.11^{+0.14}_{-0.21}$	$0.91^{+0.24}_{-0.25}$	0.51 ± 0.05	41	30
2136+141	1991 May 9	147 ± 16	4076	20.93	...	$0.50^{+0.50}_{-0.70}$	0.12 ± 0.02	22	20
2201+315	1993 May 12	553 ± 25	3329	21.09	$21.29^{+0.11}_{-0.14}$	$2.00^{+0.90}_{-0.70}$	$3.78^{+0.62}_{-0.58}$	14	32

^afrom Brunner *et al.*^b N_{H} is atomic and molecular^cfrom Zhang *et al.*




TABLE 8
SPECTRAL DISSECTION PARAMETERS

Source (1)	ν_n (2)	ν_m (3)	F_m (4)	α_{mm} (5)	α_{thick} (6)
0016+731	60	54	1.94	1.36	0.33
0133+476	42	37	3.80	0.65	1.13
0212+735	64	58	1.99	1.57	0.0
NRAO 140	78	69	2.92	1.35	0.38
3C 111	79	73	4.28	0.75	0.81
3C 120	67	67	2.63	0.80	0.41
0552+398	39	47	4.35	1.06	0.0
0642+449	75	65	4.09	1.81	0.0
0736+017	58	78	1.95	0.84	0.12
0804+499	45	57	2.15	0.63	0.56
0836+710	27	33	1.34	0.90	0.21
0923+392	...	8	4.5	0.52	...
0945+408	33	33	1.73	0.72	0.0
0954+556	19	28	0.88	0.65	0.28
1611+343	47	51	3.18	0.65	0.88
1633+382	74	94	1.82	0.60	0.56
3C 345	48	53	8.98	0.60	0.90
1954+513	27	38	2.52	0.65	0.35
2005+403	37	26	4.18	0.92	1.58
2037+511	31	50	3.88	0.63	0.11
2136+141	19	15	1.63	0.77	0.27
2201+315	26	21	4.32	0.70	1.47






TABLE 9
COMPTON CALCULATION PARAMETERS

Source (1)	z (2)	ν_m (3)	F_m (4)	α_{mm} (5)	θ (6)	$F_{\nu x, pred}$ (7)	$F_{\nu x, obs}$ (8)	δ_{min} (9)
0016+731	1.76	54	1.94	1.36	0.67	5.1×10^{-8}	0.06	0.13
0133+476	0.86	37	3.80	0.65	$\lesssim 0.08$	$\gtrsim 2.6 \times 10^5$	0.27	14
0212+735	2.37	58	1.99	1.57	0.36	7.3×10^{-7}	0.20	0.17
NRAO 140	1.26	69	2.92	1.35	0.52	4.3×10^{-7}	1.59	0.11
3C 111	0.05	73	4.28	0.75	0.50	1.0×10^{-5}	12.3	0.08
3C 120	0.03	67	2.63	0.80	$\lesssim 0.46$	$\gtrsim 1.5 \times 10^{-6}$	18.4	0.06
0552+398	2.3	47	4.35	1.06	0.47	4.6×10^{-2}	0.21	0.79
0642+449	3.40	65	4.09	1.81	0.32	0.03	0.06	0.89
0736+017	0.19	58	1.74	0.84	$\lesssim 0.35$	$\gtrsim 8.4 \times 10^{-6}$	0.50	0.15
0804+499	1.43	45	2.49	0.63	$\lesssim 0.21$	$\gtrsim 8.28$	0.24	2.0
0836+710	2.17	27	1.80	0.90	$\lesssim 0.17$	$\gtrsim 7.1 \times 10^2$	1.2	3.0
0923+392	0.70	8	4.5	0.52	0.54	2.2×10^3	0.52	5.2
0945+408	1.25	54	1.28	0.72	0.58	3.0×10^{-6}	0.09	0.15
0955+476	1.87	65	0.70	0.97	0.32	2.1×10^{-6}	0.21	0.14
1611+343	1.40	45	3.17	0.65	0.27	2.8	0.15	1.7
1633+382	1.81	94	1.23	0.6	0.33	6.7×10^{-5}	0.41	0.19
3C 345	0.60	48	9.33	0.6	$\lesssim 0.31$	$\gtrsim 24$	0.51	2.1
2136+141	2.45	15	1.63	0.77	$\lesssim 0.42$	$\gtrsim 18$	0.12	2.5
2201+315	0.30	21	4.10	0.70	0.41	2.0	3.78	0.89











TABLE 12
COMPARISON OF SPECTRAL INDICES


Source (1)	α_{mm} (2)	α_x (3)
0016+731	1.36	1.00 ^a
0133+376	0.65	$0.66^{+0.64}_{-0.36}$
0212+735	1.57	-0.60 ± 0.40
NRAO 140	1.35	$1.01^{+0.16}_{-0.17}$
3C 111	0.75	$2.25^{+0.65}_{-0.40}$
3C 120	0.80	$2.13^{+0.09}_{-0.11}$
0552+398	1.06	$0.09^{+0.36}_{-0.16}$
0642+449	1.81	$0.79^{+0.86}_{-0.59}$
0736+710	0.84	$1.24^{+0.36}_{-0.14}$
0804+499	0.62	$1.09^{+0.71}_{-0.79}$
0820+560	0.90	$0.88^{+0.13}_{-0.13}$
0836+710	0.90	0.57 ± 0.11
0917+449	0.67	$0.94^{+0.30}_{-0.24}$
4C 39.25	0.74	$1.17^{+0.38}_{-0.34}$
0945+408	0.72	$1.06^{+0.22}_{-0.24}$
0954+556	0.65	$1.30^{+0.45}_{-0.30}$
0955+476	0.97	$0.19^{+0.15}_{-0.20}$
1150+812	0.72	$0.41^{+0.19}_{-0.29}$
1611+343	0.65	1.34 ± 0.30
1633+382	0.60	$0.76^{+0.17}_{-0.20}$
1638+398	0.66	1.89 ± 0.30
1641+399	0.60	$0.90^{+0.24}_{-0.25}$
2136+141	0.77	$0.50^{+0.50}_{-0.70}$
2201+315	0.54	$2.00^{+0.90}_{-0.70}$


^aassumed value for spectral index











Multiwaveband Observations of Quasars with Flat Radio Spectra and Strong Millimeter Wave Emission

Steven D. Bloom ^{1,2}, Alan P. Marscher, E. M. Moore ³

Department of Astronomy, Boston University, 725 Commonwealth Avenue, Boston, MA
02215

Walter Gear ⁴

Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ , Scotland, UK

Harri Teräsranta, Esko Valtaoja

Metsahovi Radio Research Station, Helsinki University of Technology, Otakaari 5A ,
SF-02150 Espoo, Finland

Hugh D. Aller and Margo F. Aller

Astronomy Department, University of Michigan, Dennison Building, Ann Arbor, MI 48109

Received _____; accepted _____

¹NAS/NRC Resident Research Associate, NASA/Goddard Space Flight Center

²Current address: Infrared Processing and Analysis Center, Jet Propulsion Laboratory
and California Institute of Technology, MS 100-22, Pasadena, CA 91125

³Current address: Sterling Software, MS 210-2, NASA Ames Research Center, Moffet
Field, CA 94035-1000

⁴Current address: Mullard Space Science Laboratory, Holmbury St. Mary, Nr Dorking,
Surrey RH5 6NT, United Kingdom

ABSTRACT

We present multiwaveband observations of a well-selected sample of 28 quasars and two radio galaxies with flat radio spectra and strong millimeter-wave emission (referred to here as FSRQ’s). The data are analyzed to determine the radio to infrared and X-ray to γ -ray properties of FSRQ’s and the relationships between them. Specifically, the synchrotron self-Compton (SSC) process is examined as a likely common radiation mechanism. For most sources, the broad band spectra are still incomplete, especially in the far-infrared and ultraviolet range. Therefore precise analysis, such as model-fitting of spectra is not usually possible. To compensate partially for this, we have taken a statistical approach, and examine the relationship between high and low energy emission by using the data set for the entire sample.

We use very long baseline interferometry (VLBI) at 8.4 and 22 GHz — higher frequencies than those of previous surveys — in conjunction with nearly simultaneous radio to submillimeter-wave observations to determine the parameters of the synchrotron spectrum and to examine the compact angular structure of a subset of sources from our sample. These parameters are used to predict the SSC X-ray flux densities. Seven of thirty sources have predicted self-Compton X-ray flux densities well above the observed flux densities obtained with the ROSAT satellite unless one assumes that the radiating plasma experiences bulk relativistic motion directed toward the observer’s line of sight. Three of these seven sources are detected at γ -ray frequencies. Model spectra show that the X-rays are consistent with the first order SSC process, with the simultaneous multiwaveband spectrum of the quasar 0836+710 obtained in 1992 March being very well fit by SSC emission from a uniform, relativistically moving source. The γ -rays are not produced via second order self-Compton

scattering, but rather by either first order self-Compton scattering or some other process.

A comparison of the ROSAT X-ray flux densities and those obtained earlier with the *Einstein Observatory* show that several FSRQ's are X-ray variables on timescales of about a decade. Several sources that were observed more than once with ROSAT also show variability on timescales of 1–2 yr, with the X-ray variability in these cases often associated with millimeter-wave variability and lower VLBI core-to-jet flux ratios. Detections at γ -ray energies also appear to be related to increases in the radio to millimeter-wave flux densities.

Statistical analysis shows that the millimeter-wave and X-ray luminosities for the sample are strongly correlated, with a linear regression slope ~ 0.6 . The peak in the distribution of X-ray to millimeter spectral indices also indicates a strong connection between millimeter-wave and X-ray emission. Particularly interesting is a correlation between X-ray to millimeter spectral index and fraction of flux density contained in the VLBI core. This tendency toward higher X-ray fluxes from sources with stronger jet emission implies that the knots in the jet are prominent sources of X-rays.

1. Introduction

It is clear that in quasars there is a strong relationship between the nonthermal emission at low frequencies and that at high frequencies. Quasars are generally strong X-ray emitters (e.g., Tananbaum et al. 1979; Ku et al. 1980), and a number have been found to be strong, hard (> 100 MeV) γ -ray sources (Fichtel et al. 1994; Thompson et al. 1995; Thompson et al. 1996). Quasars with flat radio spectra — which indicate compact, milliarcsecond-scale structure — and strong millimeter-wave emission (here abbreviated as FSRQ’s) have been demonstrated to be more luminous in soft X-rays (for a given optical luminosity) than either their radio-quiet counterparts or radio sources with the same radio luminosity but steeper radio spectral index (Owen et al. 1981; Zamorani et al. 1981; Ledden et al. 1985 ; Kembhavi et al. 1986; Worrall et al. 1987). Browne et al. 1987 have also shown that for a given radio luminosity in extended (arcseconds or greater) structure, the X-ray luminosity tends to be more than an order of magnitude larger for sources with higher compact radio luminosity. Furthermore, except for the Large Magellanic Cloud, bright extragalactic hard γ -ray sources are all FSRQ’s (or BL Lac objects, which have similar radio–infrared spectra).

These strong correlations between radio and high-energy emission are readily explained if the nonthermal emission at all wavebands is produced via synchrotron self-Compton (SSC) scattering, as has been suggested by many authors (e.g., Jones et al. 1974a). This is, however, not the only possibility, since other radiative processes are likely to operate in an environment that produces the highly relativistic electrons and strong magnetic field required for SSC emission (eg., Dermer et al. 1993).

If the SSC process dominates the nonthermal emission, one would expect a correlation between X-ray and γ -ray fluxes and VLBI “core” fluxes. However, a statistical study of the (non-simultaneous) VLBI core emission and X-ray emission from compact radio loud quasars and active galaxies (Bloom & Marscher 1991) did not conclusively demonstrate a

strong correlation. Though Zhou *et al.* (1997) do find a strong relationship between high energy γ -ray fluxes and VLBI fluxes. Nevertheless, detailed studies of individual sources, both with VLBI and X-ray data, show that the SSC process is a plausible explanation for the X-ray emission (Marscher 1988; Unwin et al. 1994; Unwin et al. 1997; Eckart et al. 1986; Eckart et al. 1987). The results of these investigations have motivated us to conduct a more detailed study of FSRQ's and the relationship between the radio–infrared and the high energy radiation. The present study not only explores the statistical relationships between high and low energy emission, but also compares theoretical SSC spectra to multiwaveband data from a set of contemporaneous measurements. Observations over a large range in frequency allow for more accurate determinations of spectral index, spectral peak frequency and flux density, and consequently a more accurate prediction of the Compton flux densities.

This study, which includes high-frequency (8.4 and 22 GHz) VLBI imaging, as well as radio through γ -ray (not all wavebands for all sources) flux density measurements, constitutes the most comprehensive set to date of multiwaveband observations of bright FSRQ's. Below we summarize the observations and data analysis. We present our own VLBI, X-ray, submillimeter-wave (submm), and near-infrared (IR) data. A more extensive presentation of previous radio–IR observations used in this work is given by Bloom et al. 1994. We follow with explanations of the procedure for spectral deconvolution and determination of spectral peak frequency and peak flux density, SSC calculations for specific sources, and comparisons of model spectra against the data. In the final sections we present a statistical analysis of the sample and a general discussion of the results.

2. Observations and Data Analysis

2.1. Source Selection

The sample is taken from an essentially complete list of radio sources stronger than 1 Jy at 5 GHz, with spectral indices between 1.5 and 4.9 GHz flatter than $\alpha = 0.5$ ($F_\nu \propto \nu^{-\alpha}$) (Steppe et al. 1988). Quasars and galaxies with 90 GHz flux densities exceeding 0.5 Jy, declinations $\geq 0^\circ$, and measured redshifts exceeding 0.03 were selected from Table III of Steppe et al. 1988 to generate a preliminary sample. This resulted in a sample that had a preponderance of sources with redshifts between 0.4 and 2.25. In order to pare the sample down to a more reasonable size, within the redshift range of 0.4 to 2.25 only sources with declinations $\geq 30^\circ$ were included in the final sample. BL Lac objects, which form a class distinct from quasars with respect to their soft X-ray emission (Worrall & Wilkes 1990) are excluded, as are optically unidentified objects. The final sample contains 30 sources — 28 quasars and two radio galaxies — and is well distributed in redshift. (One source that would have been in the final sample, 3C 216, was dropped from the study due to its proximity to the Sun during the time of the submillimeter observations.) Because of telescope time allocation limitations, it was not possible to observe the entire sample at all wavelengths. Approximately one half of this sample was observed from radio to X-ray wavelengths. This subsample was essentially determined by which sources were observed (in priority order) with ROSAT prior to the shutdown of the PSPC detector. This might then have caused a selection of objects that were *a priori* deemed to be more interesting than others. Both radio galaxies, 3C 111 and 3C 120, are included in the subsample. The list of sources, along with a summary of the observations at all wavebands, is presented in Table 1.

2.2. VLBI Data

A subsample of sources has been observed with VLBI at frequencies of 8.4 and 22 GHz. For this work, seven observing sessions were conducted with either the Global VLBI

Network or the Very Long Baseline Array ¹(VLBA; after 1994). The stations of each array (defined by the epoch of the observations) are listed in Table 2. Sessions 1-6 used the Mark II VLBI recording system with a 1.8 MHz bandwidth, while sessions 7-9 used the VLBA recording system with 16 and 32 MHz bandwidth respectively. Due to the large sample size, we observed in “snapshot” mode with each source typically observed for 4 to 5 scans of one-half hour each during the allotted observing time. Typically three to six sources were observed during each VLBI session. These scan times were chosen to optimize u - v coverage. The video tapes from the individual antennas were correlated using the JPL/Caltech Block II correlator for sessions 1-6 and the VLBA correlator for sessions 7-9. All post-processing was performed at Boston University using the NRAO AIPS, the CalTech VLB software package, and the difference mapping software Difmap. The phase time and frequency derivatives (fringe delay and fringe rate) were determined using the global fringe fitting algorithm FRING in AIPS (Schwab & Cotton 1983). The amplitudes were calibrated in the usual manner as described by Cohen *et al* (1975). Inconsistencies in the calibration of flux densities were corrected by using u - v crossing points and then finding the best-fit flux density using a χ^2 test. A hybrid mapping procedure was used to obtain images of the sources (see, e.g., Pearson & Readhead 1984). Hogbom’s CLEAN algorithm was used to deconvolve the dirty beam from the dirty image to determine the true source brightness distribution. A Gaussian restoring beam with dimensions equal to those of the central portion of the dirty beam was convolved with the CLEAN components to produce a final map.

Table 3 summarizes the parameters of the final image for each source at each frequency:

¹The VLBA is an instrument of the National Radio Astronomy Observatory, a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

elliptical Gaussian restoring beam ($a_{\text{beam}}, b_{\text{beam}}$), the position angle of the beam (θ_{beam} , as measured north through east), the peak flux density per unit beam area, and the contour levels. Table 4 summarizes the Gaussian model parameters used in conjunction with the hybrid mapping procedures, as described above. These are from the model fits to the self-calibrated data. As with the Pearson & Readhead (1988) survey, the limited data only weakly constrain some model components. In Tables 3 and 4, r is the angular distance from the map center, θ_{pos} is the angular position of the component relative to the (arbitrarily placed) origin (measured from north through east), a and b are the FWHM major and minor axes of the assumed elliptical Gaussian brightness distribution of the component, and ϕ is the position angle of the major axis. The uncertainties in the flux densities given are the estimated calibration errors (as opposed to the uncertainty in the model, which is usually larger). Though we have not, in general, estimated the uncertainties for every parameter of each component, we have performed an uncertainty analysis on the angular sizes (characterized as major and minor axes of an elliptical Gaussian here) of some sources. This will be helpful later (§3.2) in determining the uncertainty of the predicted Compton flux, which is highly sensitive to the angular size value. In determining the magnitude of the $1\text{-}\sigma$ uncertainties, we use the technique of Biretta *et al* (1986). Due to the large range in the data quality of our snapshot images, there is also a fairly large range in the uncertainties of derived angular sizes. For the 4 objects observed with the VLBA and also later used for the Compton calculations, the uncertainties are 13% for 0945+408, 0955+476, 1611+343, and 2201+315 respectively.; however, for sources observed earlier with the Global VLBI Network, the typical model uncertainties are several times larger. Figure 1 displays the VLBI images at 8.4 and 22.2 GHz, in order of increasing right ascension for sources mapped with Global VLBI. Figure 2 displays the maps from the VLBA. Note the differences in angular scale, especially between most of the 22 GHz and 8.4 GHz maps.

2.3. Submillimeter and Infrared Data

A detailed description of the data analysis at submillimeter and infrared wavebands is given in Bloom *et al* (1994). In Tables 5 and 6 we present data more recently acquired with the JCMT and UKIRT. The last column in Table 5 is a measure of variability amplitude, as determined by the parameter $\frac{F_{\max}-F_{\min}}{F_{\max}+F_{\min}}$. The two measured flux densities for this calculation, extracted from the data presented here or in Bloom *et al* (1994), are from the two observations that are closest in time. This measurement of the variability amplitude is considered to be significant if there is at least a $2\text{-}\sigma$ separation between the error bars of the highest and lowest point. We have also examined the possibility of variability in the older data (between 1991 and early 1993). The sources found to be variable are discussed in §2.6.

2.4. X-ray Data

The Position Sensitive Proportional Counter (PSPC) was used in conjunction with the Woltjer Type I nested mirror system onboard the Roentgensattelit (ROSAT) to obtain fluxes over the energy range 0.1–2.5 keV. After initial processing at Goddard Space Flight Center (GSFC), the X-ray data were extracted from tape using the Post Reduction Off-line Software (PROS) routines operating under the NOAO IRAF data analysis package. Before conducting a spectral analysis, the background counts were determined for each source by calculating the number of photons within concentric annuli, centered on the source. The precise sizes of the annuli (typically $2'$ inner and $3'$ outer) were determined for each source by displaying the photons/pixel with SAOIMAGE (see Wilkes 1992). All of the objects in this sample are unresolved with the PSPC, which has a spatial resolution of about $15''$. Therefore, spatial analysis of the photon counts (i.e., mapping the source) is not possible.

The spectral analysis was conducted using the standard routines within PROS. This

consisted of modeling the data in terms of a power-law spectrum over the energy range 0.1–2.5 keV, with absorption at low energies due to intervening gas, parameterized as the column density of hydrogen atoms under the assumption of cosmic abundances. The X-ray photon spectrum is represented by the following equation:

$$\frac{dN}{dE} = K E^{-(\alpha+1)} e^{-N_H \sigma(E)} \quad \text{photons s}^{-1} \text{cm}^{-2} \text{keV}^{-1}. \quad (2.4.1)$$

The predicted spectrum (photon counts per energy bin) using this model was then compared to the observed spectrum using a χ^2 test to determine “goodness-of-fit.” The model parameters that minimize χ^2 were adopted to calculate the total flux over a specified energy band. The default values for the spectral response matrix and effective area of the detector for routines in PROS 2.0 were used to determine the source spectrum from the raw photon spectrum. Some care has to be taken, as χ^2 can be a minimum for parameter values that are physically unrealistic, such as an extremely high column density, or a photon index that is much too steep. Uncertainties in model parameters and derived flux values were determined by increasing the value of the parameter by the amount that increases χ^2 to the desired confidence level. Alternatively, in some cases, the normalization parameter, K , of equation (2.4.1) was plotted against spectral index, and the uncertainty in this parameter was taken to be the range of the 68% confidence contour. The flux density was then calculated by using equation (2.4.1) corrected for absorption, and then converting to Janskys.

Table 7 summarizes the ROSAT X-ray observations. Column (5) is the logarithm of the Galactic hydrogen column density, $N_{H\text{gal}}$. Unless otherwise noted, this is the value from the Stark *et al.* (1992) 21-cm emission line observations. However, for two sources, 3C 111 and NRAO 140, we have also added the column density (in atoms cm^{-2}) of molecular hydrogen as inferred from observations of Galactic CO (Bania et al 1991). Column (6) gives the best fit value of N_H as determined from the χ^2 minimization routine. Column (7) is the best-fit spectral index; if no uncertainties are given, then it is an assumed value.

Column (8) is the 1 keV flux density. Columns (9) and (10) give the χ^2 and degrees of freedom (D.O.F., one less than the number of parameters varied subtracted from the total number of energy channels), respectively. When N_H is determined from a best fit, then the uncertainties quoted in the spectral index and flux densities correspond to $\chi^2_{min} + 3.53$ (68% confidence for three interesting parameters). If N_H is assumed to be the Galactic value, then the subsequent uncertainties correspond to $\chi^2_{min} + 2.3$ (68% confidence for two interesting parameters). If both α and N_H are assumed then the uncertainties are for $\chi^2_{min} + 1.0$ (68% confidence for one interesting parameter). Generally, the fit is considered reasonable if $\chi^2 \lesssim \text{D.O.F.}$ Note that by this criterion two fits are not particularly good (see Table 7). The fit for the first observation of 1150+812 is poor probably because the fit was performed on the combined counts from two scans that were a month apart, each having a low number of individual counts. Also, the actual N_H value could be smaller or larger than the assumed Galactic value (see below for a discussion of this point). It is also possible that a single power-law model is not adequate for describing this source. The first observation of 1611+343 is poorly fit. Probably only the last reason is applicable here, since there are enough counts to determine the spectrum rather well, and the best-fit N_H is fairly close to the Galactic value.

In many cases the best-fit values for N_H are either considerably smaller or larger than the measured Galactic N_H . There are several likely explanations for this. If the best-fit value is larger, as is the case for 3C 120 and 0736+017, then there could be significant photoelectric absorption within the source. There could also be significant amounts of as yet unobserved Galactic molecular gas in the direction of the FSRQ. Also possible is a spectral flattening at low X-ray energies. A closer look at the photon spectrum of 3C 120 shows that the peak is poorly fit by the model, indicating that a single power-law is probably not the best model for this source. It is also possible that the Galactic column density of hydrogen is higher (or lower) in the immediate vicinity of the FSRQ as compared to the

average value over the much larger regions measured with the 2° FWHM beam of the Stark *et al.*(1992) survey. This seems to be the case for 3C 345, for which N_H measured with the smaller beam in the Elvis *et al.* (1989) study is 7.5×10^{19} , which is certainly closer to the best-fit value of N_H for the 1990 observation than is the value $N_{H_{\text{gal}}} = 3.6 \times 10^{20} \text{cm}^{-2}$ derived from Stark *et al* (1992). These possibilities have been discussed in more detail by Wilkes & Elvis (1987) and Worrall & Wilkes (1990), among others. However, the higher resolution H I observations of Elvis *et al.* (1989) show that $N_{H_{\text{gal}}}$ is smaller than derived from Stark *et al* (1992) for 0736+017 and only 2% larger for 3C120. It is still possible that $N_{H_{\text{gal}}}$ in the immediate direction of the quasar was larger at the time of the X-ray observations, since temporal variability of $N_{H_{\text{gal}}}$ cannot be excluded (see, e.g., Fichtel et al. 1994). For 0736+017, the inferred column density from CO observations (Liszt et al. 1993) is $5.6 (\pm 0.8) \times 10^{20} \text{cm}^{-2}$, which can account for the difference. This also results in a much steeper derived value of the spectral index, which is more in agreement with other low-redshift active galaxies and quasars (see discussion below). If the best-fit value of N_H is smaller than the Galactic value, as is the case for 0133+476, 3C 111, 1633+382, and 3C 345, then it is possible that these sources are not adequately described by a single power-law. The most likely possibility is a soft X-ray excess, perhaps from thermal black body X-ray emission (from an accretion disk) with a steep slope.

An analysis for some sources in this sample has appeared in other summaries of X-ray emission from blazars (see Comastri et al. 1997; Sambruna 1997) . Most of the parameter determinations (such as spectral index and column density of hydrogen) are very similar. Our flux density values do occasionally differ, mainly because we base our flux values on best fit parameters, whereas Sambruna (1997) use fixed $N_{H_{\text{gal}}}$ to calculate flux densities. In addition, their source/background counts determination differs from ours slightly. The first observation of 1611+343 is the only case for which the flux densities differ by more than 5% (we present a value which is 50% lower).

2.5. Gamma-Ray Data

Several sources in our sample were detected with the EGRET instrument on the *Compton Gamma-Ray Observatory*. The fluxes for EGRET detected objects can be found in Thompson *et al.* (1995) and upper limits for some sources in this sample can be found in Fichtel *et al.* (1994). The dates listed in Table 1 are for the EGRET Viewing Periods, covering the area of the source, which were closest in time to the other multi-wavelength observations. Calibration of the instrument and determination of fluxes are discussed in Hartman *et al.* (1992), Mattox *et al.* (1993), and Mattox *et al.* (1996), respectively.

2.6. Summary of Source Properties

Due to incomplete time coverage, it is not possible to analyze thoroughly the variability properties of each source. However, we can discuss some general results. Of the 17 sources with repeated millimeter/submillimeter wave observations reported here and in Bloom *et al.* (1994), ten (0133+476, NRAO 140, 3C 111, 0642+449, 0736+017, 0836+710, 3C 345, 2005+403, 2037+511, and 2201+315) are significantly variable at one or more wavelengths. Six of these (0133+476, NRAO 140, 0736+017, 0836+710, 3C 345, and 2201+315) are significantly variable at all measured wavelengths. The timescales are roughly 7 to 14 months; no source has a variability amplitude parameter that exceeds 0.5.

With the X-ray observations presented earlier, combined with observations by previous spacecraft missions (HEAO A-1, Einstein, EXOSAT, and Ginga), we have determined the crude X-ray variability characteristics of the sources in the sample, using the same criterion for significance as for the millimeter observations. We note that in some cases it is difficult to establish variability, especially if the observations being compared were made with different instruments or very different spectral fit parameters (as mentioned above).

The sources appear to fall into several general categories (Malaguti et al. 1994; Bloom 1994 for references to the published X-ray data for each source). Of the 24 sources in our sample with ROSAT X-ray observations, seven (0133+476, 0552+398, 0820+560, 0917+499, 0954+556, 0955+476, and 1638+398) have indeterminate X-ray variability due to insufficient observations. For an additional seven sources (0016+731, 0642+449, 0736+017, 0804+499, 1150+812, 2136+141, and 2201+315) variability was not detected, though some of these are low amplitude variables at mm wavelengths (e.g., 0736+017; see Tornikoski et al. 1994). Two sources, 0945+408 and 1633+382, are observed to be significantly variable in X-rays (and, for 1633+382, γ -rays; see Thompson et al. 1995) only, whereas three additional sources (NRAO 140, 0836+710 and 4C 39.25) show possible X-ray variability that is temporally correlated with radio or mm-wave variability. With each of these three sources, the mm flux varied by the same factor as the X-ray flux (see Marscher 1988 for NRAO 140; this work and Bloom *et al.* (1994) for 0836+710 and 4C 39.25). These last five sources mentioned were all variable with constant X-ray spectral index, which is typical of nonthermal processes. Five additional sources (0212+735, 3C 111, 3C 120, 1611+343, and 3C 345) had complex variability properties, such as variation in spectral index as well as intensity. All of the sources with $\lesssim 80$ % of the VLBI flux density in the core at 8.4 GHz are X-ray variable, falling in one of the last three categories. The possible relationship between X-ray and milliarcsecond-scale structure is discussed in §4.2 .

3. Spectral Analysis

3.1. Spectral Deconvolution

In this section, the radio to submillimeter and infrared spectra of Bloom *et al.* (1994) are used to determine the synchrotron spectral turnover frequencies for each source in the sample with sufficient data. In addition, for sources observed with both ROSAT and VLBI,

X-ray measurements are compared with the theoretical spectra of synchrotron self-Compton (SSC) models generated by the techniques described below. For sources with published EGRET γ -ray data, those data are also included for further comparison with the theoretical spectra. The total flux density spectra presented in Bloom *et al.* (1994) are dissected into the spectra of individual VLBI scale components using the total flux density data and flux densities from the best-fit models to the 8.4 and 22 GHz VLBI visibilities. The emphasis of the spectral decomposition is on finding the spectral turnover frequency and flux density of the bright “VLBI core” component usually located at one end of a source. Due to the limited data and the weakness of the secondary components we do not attempt to fit the spectra of any non-core “knots” (except for the peculiar source 4C 39.25, which was analyzed by Zhang & Marscher 1994 as part of a separate study).

In Figure 3 the spectral dissection for a representative source is shown. The turnover frequency ν_m is determined from the data by first calculating the frequency ν_n at which the optically thin and optically thick spectra (of slope α_{mm} and α_{thick} , respectively) intersect (Jones et al. 1974b ; Marscher 1977). In some cases this is not straightforward, since the VLBI spectra are complex (e.g., 3C 120 and 1633+382). For such sources, there is significant blending of components at 8.4 GHz that are separated in the higher resolution 22 GHz maps. There are also additional components not visible in the 22 GHz images because of over-resolution and/or steep spectra. There are insufficient data to separate these components properly, thus only the 22 GHz core flux is used with the total flux density data to determine the spectral turnover. In sources with no VLBI data, best estimates are made using the total flux density data. We also note that in some cases, the VLA and/or VLBI data points exceed the values for the single-dish measurements (Figure 3). Since it is obviously not possible for the small scale components to have greater flux density than the entire source seen by a single dish, these differences in flux are attributable to differences in calibration. Since all of the sources have falling millimeter and submillimeter spectra,

there is a firm upper limit of 100-150 GHz to the turnover frequency. All sources in the sample have optically thick spectral indices below the turnover that are much flatter than the value of 2.5 expected from a homogeneous source. These flatter indices are indicative of sources with decreasing radial gradients in magnetic field B and electron energy distribution normalization factor N_0 . Using the expressions of Marscher (1977), who considered such cases, the actual synchrotron turnover frequency ν_m is related to the intersection value ν_n by a constant factor, which in most cases is close to unity. The quantity F_m is the turnover flux density determined by extrapolating the optically thin spectrum down to frequency ν_m . The values of ν_n , ν_m , F_m , α_{mm} , and α_{thick} for each source are listed in Table 8.

For sources observed with VLBI, the angular sizes are derived from those obtained from the models that best fit the visibilities as described in §2.2. Though elliptical Gaussians are used to represent the brightness distribution of each source component, for the SSC calculations the visibilities can equivalently be represented by a uniform sphere with angular diameter $\theta_{us} = 1.8\sqrt{\theta_a\theta_b}$ (Marscher 1987). The equivalent size for a spherically symmetric, non-uniform source is $(1.8 k_\theta)^{-1}\theta_{us}$, where k_θ is a parameter in the range 1–2 (Marscher 1977). The value θ_{us} is used in the subsequent modeling calculations. This angular size value is only strictly valid if the core is known to be optically thin. In most cases, our spectra reveal the core to be optically thick; however, we use these optically thin values for consistency with the following Compton calculations (which assume the source is optically thin).

3.2. Synchrotron self-Compton Calculations

Once F_m and ν_m are determined, these values as well as millimeter spectral indices and angular sizes are used with the following equation (multiplied by a constant factor; see

Marscher 1987):

$$F_{\nu}^{1C} \propto \theta^{-2(2\alpha_{mm}+3)} F_m^{2(\alpha_{mm}+2)} \nu_m^{-3\alpha_{mm}+5} E_{keV}^{-\alpha_{mm}} \ln(\nu_2/\nu_m) \left(\frac{1+z}{\delta} \right)^{2(\alpha_{mm}+2)}. \quad (3.2.1)$$

Here, ν_2 is the upper cutoff frequency to the synchrotron spectrum and δ is the Doppler factor corresponding to bulk relativistic motion. We use eq. (3.2.1) to predict the X-ray flux density at 1 keV under the initial assumption that there is no relativistic beaming ($\delta = 1$). The angular sizes used in this equation should be measured at the turnover frequency. In general this is not the case, since VLBI observations were only undertaken at 8.4 and 22 GHz, below ν_m for most sources. In the case of a turnover frequency $\nu_m > 22$ GHz, the measured angular size is expected to follow the inequality $\theta(\nu) > \theta(\nu_M)$, and the Compton flux is underestimated. If the source is transparent at 22 GHz, then the measured angular size is acceptable. For sources with IR data, the cutoff frequency ν_2 can be estimated, and is typically $\sim 10^{14}$ Hz. Otherwise, this frequency is assumed to be 10^{14} Hz.

In a number of cases the X-ray flux densities predicted in this way are much greater than those measured. If relativistic beaming is taken into account, a minimum Doppler factor can be derived from the measured and predicted X-ray flux densities:

$$\delta_{min} = \left(\frac{F_{\nu x, obs}}{F_{\nu x, pred}} \right)^{-\frac{1}{2(\alpha_{mm}+2)}}. \quad (3.2.2)$$

The best estimate of the Doppler factor is then used with the initial estimate of the X-ray flux density to generate model first and second order SSC spectra. Model spectra for selected FRSQ's with minimum Doppler factors obtained from the SSC calculations are shown in Figures 4 (0133+476) and 5 (0836+710). The parameters used to generate the models are listed in Table 9. The SSC calculations show that seven of sixteen sources observed with both VLBI and ROSAT have predicted X-ray flux densities at 1 keV greater than the observed values. The resolution of this discrepancy requires relativistic beaming of the radiation.

To evaluate more clearly what these predictions mean, we can roughly estimate the uncertainty in the SSC predicted flux and the Doppler factor using standard propagation of errors techniques (i.e., Bevington 1969). Though this strictly only holds for parameters which have normal distributions and symmetric errors, a meaningful result for the uncertainty can be evaluated if we treat the terms of eqs. 3.2.1 and 3.2.2 logarithmically:

$$\sigma_{tot}^2 = a^2 \sigma_{\log\theta}^2 + b^2 \sigma_{\log F}^2 + c^2 \sigma_{\log\nu}^2 + \dots \quad (3.2.3)$$

Here, σ_{tot} refers to the total logarithmic uncertainty in the predicted Compton flux. a , b , c refer to the powers involving spectral index which appear in eq. (3.2.1). The corresponding subscripts refer to angular size, maximum synchrotron flux density and the spectral turnover frequency. There are then other smaller or unquantifiable terms which we have not written down (but we discuss them below). Following this technique, and assuming approximately 50 % errors in the angular size, and 40% errors in the turnover flux and frequencies, and a spectral index of approximately 0.7, the uncertainty in $\log F_{\nu x, pred}$ is about 2. Realistically, since the spectral index itself can be uncertain by a few percent, and since there are very likely to be correlated error terms which we haven't included, this number could be closer to 3. Though correlated error terms could also reduce the overall uncertainty, in the absence of more information we feel that it is wise to adopt the larger number. This number corresponds to a “worst case scenario”. As discussed earlier in §2.2, some of the VLBI data have angular sizes with uncertainty as small as 13%. The resulting uncertainty would be smaller by more than an order of magnitude in those cases (~ 50). However, in general, over-estimations of the Compton flux, are only significant if this is an over-evaluation by more than 3 orders of magnitude. By this standard, all of the over-estimates in Table 9, except for 0133+476, are marginal, however many of the underestimates are not. Using similar analysis on δ_{min} , and realizing the $F_{\nu x, pred}$ is the main contribution to the uncertainty, we calculate that the uncertainty in the log of the minimum Doppler factor is about 0.3. Thus, we conclude that the uncertainties are

too large, generally, to determine whether Compton scattering and Doppler beaming are necessary to reconcile predictions with the X-ray data.

In some cases the X-ray spectral slopes predicted by the SSC mechanism disagree with the slopes determined from fits to the ROSAT data. This could be due to non-SSC contributions (e.g., from a “soft X-ray excess” component) or simply a poorly derived X-ray spectrum arising from the limited energy range to which the ROSAT PSPC is sensitive. Nonuniform models (e.g., shock waves in which the electrons suffer radiative losses; Marscher & Gear 1985) can produce self-Compton X-ray spectra that are flatter than the optically thin synchrotron spectra.

For 0836+710, the γ -ray spectrum of 1992 March is roughly in agreement with the predicted composite first and second order multiwaveband spectrum (Figure 5). These results are similar, in general, but not in detail, with the modeling results for 0836+710 presented by Comastri *et al.* (1997). The model predicts that the source should have become faint at γ -ray energies by 1992 Oct-Nov, and indeed no detection was reported when the source was next observed in early 1993 (Thompson et al. 1995). This illustrates the value of contemporaneous multiwaveband observations in testing models: source variability invalidates tests based on non-contemporaneous measurements.

In addition to modeling the spectra, we use observed synchrotron parameters, such as F_m , ν_m , α , ν_2 , and θ , in Table 10 to determine physical parameters such as N_0 and B . These results show that the magnetic field B ranges from 1×10^{-2} to 140 (δ_{min}/δ) G and N_0 from 9×10^{-13} to $2 \times 10^{-3}(\delta/\delta_{min})^{2(\alpha+2)}$ (cgs units).

For sources previously observed to exhibit superluminal motion, the range of possible values for the Doppler factor derived from the apparent transverse velocities v_{app} can be compared to those derived using eq. (4.2.2). Known superluminal sources with measured X-ray fluxes lower than the predicted values tend to have minimum required values of δ

that are compatible with the observed values of v_{app}/c , as should normally be the case: 0836+710 ($v_{\text{app}}/c = 15h^{-1}$ [Krichbaum et al. 1990], $\delta > 3$), 4C 39.25 ($v_{\text{app}}/c = 3.5h^{-1}$, $\delta > 5.2$ [Marscher et al. 1991], and 3C 345 ($v_{\text{app}}/c = 8h^{-1}$, $\delta > 2.1$; see Unwin et al. 1994). (Here we adopt the values $q_0=0$ and $H_0 = 100h \text{ km s}^{-1} \text{ Mpc}^{-1}$.) However, this need not be the case for all sources, since the value of δ can be high while v_{app}/c is low for jets pointing essentially along the line of sight. In such cases, the jet should not appear as extended as in the high v_{app}/c sources.

4. Statistical Analysis

We use correlation and linear regression to test for and to model approximately the bivariate relationships among the data. We employ these techniques with emphasis on those parameters most likely to be related to X-ray emission and γ -ray emission. Since the SSC process predicts a specific relationship between high energy flux densities, radio flux density, spectral turnover frequency, and angular size, these parameters are studied most closely. In addition, other parameters such as the X-ray and millimeter-wave spectral indices, and VLBI compactness, are examined.

All of the luminosities in this section are calculated assuming $q_0=0$ and $H_0 = 100h \text{ km s}^{-1} \text{ Mpc}^{-1}$, with h arbitrarily chosen to be 0.75.

4.1. Survival Analysis—Correlation and Regression

Due to the presence of upper and lower limits to some of the measured or inferred quantities (data with limits are often referred to as “censored” data), survival analysis techniques are used to evaluate more accurately correlation coefficients (with their associated probabilities) and regression slopes (Isobe et al. 1986). In general, these techniques take

into account censored data by assuming a form for the distribution of the censored values relative to the measured ones. The ASURV Rev 1.1 statistical software package is used to calculate correlation properties for censored data (Isobe & Feigelson 1990). As pointed out by Isobe et al. 1986, spurious correlations (or spurious null correlations) can occur if these limits are disregarded.

There are many correlation tests and linear regression methods. In general, since they are both non-parametric, the Spearman rank correlation coefficient and Kendall’s τ are best suited to test for correlation among two variables for which the nature of the parent distributions is unknown. The Spearman rank coefficients are provided in the summary of our results in Table 11.

The Buckley-James method (BJ) is used for determining regression coefficients. This method is preferred over other methods, since it does not assume a Gaussian distribution of the residuals about the regression line. In cases for which there are limits present in the independent variable (e.g., the possible dependence of X-ray flux density on angular size), only Schmitt’s regression method (SB) can be used. However, this method is not considered preferable for general use, since it involves arbitrary binning of the data. (For a more detailed discussion of the relative merits of these methods see Isobe et al. 1986). When no limits are present, the regression results are nearly identical to the results from standard methods (e.g., Bevington 1969).

Since several previous studies have shown a strong correlation between X-ray and radio luminosities for radio loud quasars (e.g., Worrall et al. 1987), we examine this relationship first. We consider separately the correlation between X-ray luminosity and radio core luminosity, X-ray and millimeter luminosity at 270 GHz, and X-ray and radio luminosity at spectral turnover. The analogous relationships between the flux densities as well as between the flux density at each waveband and redshift are studied, to ascertain whether

the luminosity correlations are intrinsic or spurious.

As can be seen in Figures 6–11 the X-ray and millimeter flux densities are correlated, the X-ray and radio core flux densities are uncorrelated, and the X-ray and turnover flux densities are weakly correlated. All of the luminosity-luminosity plots in these figures show moderate or strong correlations. The X-ray/millimeter relationship may be stronger than that of the X-ray/core because the core is optically thick. Optically thick synchrotron emission only makes a very small contribution to the total soft radiation field which is upscattered to X-ray energies (Band & Grindlay 1986). A summary of the correlation and regression analysis is given in Table 11. Regression results are given only if the probability that a correlation is not present is less than 20%.

If the correlation between the luminosities were due solely to the common distance factors used to derive them, then the regression slope would be close to unity. However, the regression slope is less than one in all cases. Isobe *et al.* (1986) show through Monte Carlo simulations of data from flux limited samples that if the underlying physical relationship between the luminosities is of the form $L_2 = KL_1^\alpha$, then luminosity diagrams will reflect this through the regression slope, as long as upper limits are included in the analysis. The resulting flux–flux plots show a correlation with considerably steeper slopes. Although the underlying physical relationship between the luminosities in this sample is likely to be more complex than the one assumed by Isobe *et al.* (1986) their results do indicate that, for some forms of luminosity functions, the data will show the intrinsic correlation in the luminosity-luminosity plots even if the sample has a distance-luminosity bias. To evaluate the possibility of spurious luminosity correlations due to common redshift factors, we have used the partial correlation methods of Padovani (1992). This analysis calculates the correlation between two variables, while effectively removing the effects of other known variables. We find that there is still a moderate correlation between L_x and both L_{mm} and

L_c ($\rho_s = 0.28$ and 0.39 respectively) , after the correlations of each luminosity with redshift have been negated. However, the partial correlation between L_x and L_m is weak, indicating that the apparent relationship between these two luminosities found earlier is probably spurious. The results are not strongly affected if the extremely low redshift objects 3C 111 and 3C 120 are removed from the sample (as is seen in Table 11).

Figures 12–17 give correlation plots and histograms of various other interesting observational parameters. Owen *et al.* (1981) used the histogram of the effective millimeter to X-ray spectral index to show that the X-ray fluxes are closely linked to the millimeter fluxes. Bloom & Marscher (1991) showed that this also holds in general for the radio cores of quasars, even though the observations were often separated by years. Figure 15 shows that this is also true for the present sample, with millimeter and X-ray observations separated by 6 months at most. Inspection of Fig. 12 and Tables 11 & 12 show that there is a weak anti-correlation between X-ray spectral index and both millimeter and infrared spectral indices. This would tend to show that a harder component to the X-ray emission is visible when the synchrtron spectrum is steep, and not contributing as much to the X-ray flux via the SSC mechanism. This finding is somewhat surprising, since we would expect that sources with less significant SSC emission would then be dominated by the softer (steeper) spectrum of an accretion disk. Though, since several of the sources with the steepest millimeter spectral indices are at high redshift, it could be that the softest X-rays from an accretion disk are redshifted out of the ROSAT band. The anti-correlations of the spectral indices are especially interesting considering that the fluxes in the millimeter and X-ray wavebands (and the luminosities) are correlated. It would be hard to argue in favor of the SSC mechanism (at least in a simple form) if these correlations are observed in the future for simultaneous data from a large set of objects. However, without detailed X-ray spectral analysis of individual sources (which will have to await AXAF) it is difficult to analyze this result much further. There is also a strong correlation between α_{mmx} and

VLBI core fraction at 8.4 GHz, as seen in Figure 16. Here, core fraction is defined to be the fraction of total milliarcsecond-scale flux density (as determined by model fitting) that is in the core component. Note that there is one outlying point, which corresponds to 3C 120. Even with 3C 120 excluded the correlation is highly significant and the regression relation remains similar to the earlier case. The correlation weakens at 22 GHz (Fig. 17). Although this may be due to the inferior calibration at high frequencies, it is also true that the jet (as opposed to core) components tend to be weaker, and the core stronger, at 22 GHz than at lower frequencies. The correlation suggests that more X-rays are created for a given mm-wave luminosity if jet components are more prominent. That sources with more prominent jet components should be brighter at high energies was also suggested by Wehrle & Mattox (1994) in their correlation between VLBI structure and γ -ray detections, although their result was of marginal statistical significance. It is certainly not expected if the bulk of the high-energy emission comes from the quiescent jet, as in the Ghissellini *et al.* (1985) model.

There are various explanations for this correlation. If the bulk Lorentz factors for each source are similar, then the sources with smaller angles between the jet axis and the line of sight will have larger Doppler beaming factors. Such sources will be more severely foreshortened as viewed by the observer, and thus the jet “knot” components will not be as easily detected (because of blending with the core) as for jets that are less highly beamed. This effect will also tend to decrease the SSC X-ray flux for a given millimeter flux (all other quantities, such as turnover frequency, being roughly equal; see eq. [3.2.1]). Recent VLBI and X-ray observations of 3C 345 (Unwin *et al.* 1997) indicate that there are states in which the X-ray emission emanates from the jet and not the nucleus, as one might normally expect. At these times the relative contribution of radio flux from the extended jet is much higher. However, we note that α_{mmx} was uncorrelated with the core fraction as this particular flare progressed in 3C 345.

Another possible explanation is that the sources with less prominent jets have wider opening angles, so knots expand rapidly, which would decrease their SSC emission. If this is the case, such sources should be found to be more highly variable (at least in the decay stages of flares) than those with prominent jets. It may also be that the sources with strong jets are in an enhanced state (e.g., from shock compression), which would increase the ratio of X-ray to millimeter flux as compared to the other sources. However, for several individual sources (NRAO 140, 0836+710, and 4C 39.25), this ratio seems to be preserved during flares. As mentioned in §2, long term X-ray variability appears to be related to 8.4 GHz VLBI structure as well, though this information is probably too limited. In the Melia & Königl (1989) model, the X-rays arise from inverse Compton scattering of seed photons from an accretion disk by relativistic electrons deep in the jet. Because of the geometry of the scattering, the X-ray emission is lower along the jet axis than at a small angle (roughly the inverse of the Lorentz factor of the jet); viewed from this angle, the jet components would have maximum visibility.

There are no significant correlations between X-ray flux density and the other parameters used to predict the X-ray flux. This is somewhat surprising since one might expect strong correlations from relations such as eq. (3.2.1). A lack of correlation between X-ray flux density and angular size can, however, be understood since many of the angular sizes are actually upper limits, and other angular sizes may represent the blending of two or more components. Furthermore, the actual angular sizes at the turnover frequency are likely to be smaller than those measured at 22 GHz (see §3). The weak correlation between X-ray flux density and turnover frequency can be understood since many of the turnover frequencies are determined with limited data and are approximate. This uncertainty would then also be exacerbated by an inaccurately determined turnover flux density. Of course, the simplest explanation for all of these non-correlations is that the X-ray flux is not SSC in origin, and thus none of the correlations would be strong. If this is the case, the actual

Doppler factors must exceed the minimum values obtained from the SSC analysis (Table 9).

4.2. Distributions

The histograms of the observed properties of FSRQ’s show which values are most common among the objects. These reveal both selection effects and the physical aspects of the objects, as is demonstrated below. The distribution of turnover frequencies (Fig. 13) shows a very broad peak at roughly 50 GHz, similar to that found by Gear *et al.*(1994). The distribution cuts off well below 150 GHz at high frequencies. If the flat spectra are caused by a jet with radial gradients in magnetic field and electron density (Marscher 1977 discusses the spherical case; see Marscher 1980; Königl 1981 for application to jets), then the angular size of the emission region, observed in the partially opaque portion of the spectrum, decreases inversely (the precise dependence is a function of the gradients in B and N) with frequency up to the spectral turnover. Thus, the spectral turnover frequencies are indicators of the innermost radius of the source. That the spectral turnovers occur at frequencies ~ 50 GHz, rather than up to an order of magnitude higher (as has been observed for some BL Lac objects and flaring quasars; see Brown et al. 1989; Gear et al. 1994) indicates that the smallest radius is farther out from the central engine than is implied for these other sources. The relationship $\theta_m \propto \theta_{22}(\nu/22GHz)^{-k_2}$ is used to approximate the size of the inner core for the sources in our sample, where θ_{22} is the angular size measured at 22 GHz. If that size is an upper limit, a value equal to half of the upper limit is used. The exponent k_2 is a constant of order unity that depends on the optically thick spectral index and is enumerated in Marscher *et al.*(1977). A histogram of this estimated “innermost radius” is provided in Figure 14. The peak in the distribution of turnover frequencies (see Fig. 13) is an indicator of the peak in the histogram of the inner radius at ~ 0.6 parsecs. Though, considering our previous discussions (§2.2; §3.1-3.2) regarding

the uncertainty of the angular size, assuming an optically thin/thick source, and possible blending of components, we warn that this number should be seen as a rough estimate of where a physical change in the jet should take place and not as a canonical number. The physical meaning of this inner radius is discussed in §5 below.

The distribution of millimeter spectral indices (see Table 12) has a sharp peak at ~ 0.7 , and is clearly skewed toward steeper spectral indices.

5. Discussion

We have shown that the high energy properties of FSRQ’s are related to their radio to infrared properties. In particular, the mm–X-ray luminosity correlation is strong, as found in previous studies. In addition, we find an anti-correlation between X-ray flux and VLBI core dominance, which suggests that the knots which comprise the observed radio jets in FSRQ’s contribute substantially to the X-ray emission.

That the distribution of spectral indices peaks near $\alpha_{mm} \approx 0.7$ indicates that most of the sources contain a distribution of relativistic electrons with a canonical $E^{-2.4}$ spectrum. However, several sources have much steeper millimeter-wave spectral indices, as expected for radiative energy losses, or emission from an inhomogenous jet (Marscher 1980). This point is further borne out by the infrared observations. Of the fifteen sources for which we have obtained infrared data (reported here in Table 6 and in Bloom et al. 1994), six show infrared spectral indices that are 0.5 to 1.0 steeper than the mm-wave spectral indices, which is again consistent with the basic predictions for radiative energy losses. However, seven sources have infrared spectral indices that are flatter by similar amounts, probably because of another (thermal?) emission component becoming prominent toward higher frequencies.

The combined peaks in the distributions of spectral turnover frequency and angular size are shown above to indicate a characteristic size of about 0.6 pc for the radio cores of the FSRQ’s observed with VLBI. In an inhomogeneous jet model with radially decreasing magnetic field and electron energy normalization, the synchrotron turnover frequency is predicted to increase with decreasing radius (Marscher 1980; Ghissellini et al. 1985). Hence, if jets extended down to some arbitrarily small radius, this would result in much higher observed turnover frequencies. Thus, there is clearly some physical inner cutoff to the radio jet. The exact nature of this is unknown. There could be a change in geometry at this radius. That is, the jet could make a transition from a conical geometry to one which has a weaker dependence on radius, such as a paraboloid (e.g. Marscher 1980). Magnetic field and electron density, and therefore the turnover frequency, would then depend less strongly on radius. This transition in geometry could be caused by a change in the external pressure on the jet or the cooling of the protons to nonrelativistic mean internal energies. A change in Doppler factor along the jet, either from a change in jet speed or orientation, could also have the effect of creating a cutoff in the emission since the observed flux from the jet follows $F_\nu \propto \delta^{3+\alpha_r}$. Millimeter-wave VLBI observations are important to study this transition region.

The lower redshift objects appear to have distinct X-ray properties. For example, both 3C 111 and 3C 120 have very steep X-ray spectra ($\alpha_x > 2$) and the X-rays could have a thermal origin (Dermer et al. 1993; Grandi et al. 1997). This steep soft X-ray spectral component would lie at frequencies below the ROSAT PSPC response at the higher redshifts of the quasars in the sample. Indeed, at higher X-ray energies both radio galaxies have flatter spectra, similar to quasars (Petre et al. 1984). Leach *et al.* (1995) have shown that the spectral slope of the low-redshift quasar 3C 273 is -2.7 in the low energy band (0.1–0.3 keV) of the ROSAT PSPC and -0.5 in the hard band (1.5–2.4 keV). Also, as discussed earlier, 0736+017, at a redshift of 0.19, has an X-ray spectral index $\gtrsim 2$ if

photoelectric absorption by Galactic molecular gas is taken into account.

Three sources, 0836+710, NRAO 140, and 4C 39.25, clearly show radio to millimeter variability that is directly proportional the X-ray variability. As discussed by Marscher (1988), this would occur if the quantity $N_0 x$ is nearly constant with time, with N_0 varying inversely as the line of sight distance through the source x . However, an increase in the Doppler factor would also have a similar signature (see Marscher et al. 1992 and below). It is further likely, from the general considerations of Bloom & Marscher (1996) and the SSC calculations in §3, that the high states of radio to X-ray (and in some cases γ -ray) emission are associated with higher high-energy cutoffs to the electron distribution. This increases the X-ray flux density by another factor of $2 \ln \gamma_{max}$. This would predict that if the γ -rays are from first order SSC, then the X-ray flux density should increase by a slightly larger factor than the synchrotron flux density. In cases for which the entire spectrum increases by a fixed amount, the increase can be attributed to a change in the Doppler factor.

Models that involve the scattering of external radiation from the relativistic electrons in jets (e.g., Melia and Königl 1989; Dermer & Schlickeiser 1993) would also explain this linear relation between millimeter and X-ray flux if changes in N_0 were solely responsible for the variations in the synchrotron and scattered fluxes. This would be the case if density fluctuations occurred while the magnetic field remained constant (see Bloom & Marscher 1996).

Of the six sources in our sample that were detected in high energy γ -rays, two were in a higher state at mm or cm wavelengths at the time of the detection. 0836+710 showed a factor of two increase in the millimeter and X-ray flux densities when it was detected in γ -rays relative to earlier times. Though 0804+499 was only observed at millimeter wavelengths several months before and after the γ -ray detection, cm-wave flux densities clearly increased by a factor of 3 between August 1991 and December 1992. The other

sources (0917+449, 0954+556, 1611+343, 1633+382) were not well observed during the time of the EGRET observations.

The physical relationships among the various parameters are often complicated and hidden by selection effects. We have used statistical tests to attempt to reveal underlying relationships. We find that the X-ray luminosity is correlated with the radio core and millimeter-wave luminosities. However, the strong correlation is probably due in part to luminosity–redshift bias.

There are many aspects of this work that should be continued to gain a better understanding of the relationship between the radio to IR and X-ray to γ -ray properties of FRSQ’s. Because many sources in this sample exhibit spectral turnovers near 40 GHz, and these turnovers indicate that the emission originates from the innermost region of the jet, they should be observed with VLBI at 43 GHz (VLBA) and in the 1-3 millimeter range (various arrays) to try to resolve these inner regions and gain a better understanding of the physical conditions at the base of the radio jet. In addition, the angular sizes derived from these VLBI images can be used for more accurate Compton calculations in conjunction with X-ray and γ -ray observations. Simultaneous multiwaveband and VLBI observations would relate the variations at different frequencies both to each other and to changes in the structure of the jet. Time-dependent models of SSC emission from relativistic jets should be developed to replace the crude uniform spherical source approximation used in this study. Finally, complete samples chosen using various well defined criteria should be studied thoroughly to establish statistical relationships as well as to elucidate the effects of those selection criteria. Together, future studies such as these should lead to a more complete understanding of the nature of the nonthermal emission in FRSQ’s.

We thank the anonymous referee for many useful comments and patience while the final manuscript was being revised. The authors gratefully acknowledge the use of data from the

service observing programs at both the JCMT and UKIRT, and thank the staffs of these telescopes for carrying out the observations on our behalf. The JCMT is operated jointly by the UK Particle Physics and Astronomy Research Council, the Netherlands Organization for the Advancement of Pure Research (ZWO), the Canadian National Research Council (NRC), and the University of Hawaii. UKIRT is operated by the Royal Observatory, Edinburgh on behalf of the UK Particle Physics and Astronomy Research Council. The National Radio Astronomy Observatory is operated by Associated Universities, Inc., under cooperative agreement with the US National Science Foundation. This material is based upon work supported by the NSF under grants AST-9116525 (Boston University) and AST-9120224 (University of Michigan), and by NASA under grants NAG5-1566, NAG5-1943 (Boston University). S. Bloom also gratefully acknowledges a grant by the Israel Council for Higher Education during his stay at Technion (Israel Institute of Technology), and an NRC Research Fellowship while at NASA/GSFC.

REFERENCES

- Band, D. L. & Grindlay, J. E. 1986, *Ap J*, 308, 576
- Bania, T. M., Marscher, A. P., & Barvainis, R. 1991, *AJ*, 101, 2147
- Bevington, P. R. 1969, *Data Reduction and Error Analysis for the Physical Sciences* (New York: McGraw-Hill), 204
- Biretta, J. A., et al. 1986, *ApJ*, 308, 93
- Bloom, S. D. 1994, Ph. D. Dissertation, Boston University
- Bloom, S. D. & Marscher, A. P. 1991, *ApJ*, 366, 16
- Bloom, S. D. & Marscher, A. P. 1996, *ApJ*, 461, 457
- Bloom, S. D., Marscher, A. P., Gear, W. K., Aller, H., Aller, M., & Teräsranta, H. 1994, *AJ*, 108, 398
- Brown, L. M. J., et al. 1989, *ApJ*, 340, 149.
- Browne, I. W. A., & Murphy, D. W. 1987, *MNRAS*, 226, 601
- Brunner, H., Lamer, G., Worrall, D. M., & Staubert, R. 1994, *A&A*, 287, 436
- Cohen, M. H., et al. 1975, *ApJ*, 201, 249
- Comastri, A. et al. 1997, *Ap J*, 480, 534
- Dermer, C. D. 1993, in *Compton Gamma-Ray Observatory*, ed. M. Friedlander, N. Gehrels, & D. Macomb (New York: AIP), 541
- Dermer, C. D. & Schlickeiser, R. 1993, *ApJ*, 416, 458
- Eckart, A., et al. 1986, *A&A*, 168, 17

- Eckart, A., et al. 1987, A&AS, 67, 121
- Elvis, M., Lockman, F. J., & Wilkes, B. J. 1989, AJ, 97, 777
- Fichtel, C. E., et al. 1994, ApJS, 94, 551
- Frail, D. A., Weisberg, J. M., Cordes, J. M., & Mathers, C. 1994, ApJ, 436, 144
- Gear, W. K., et al. 1994, MNRAS, 267, 167
- Ghissellini, G., Maraschi, L., & Treves, A. 1985, A&A, 146, 204
- Grandi, P. et al. 1997, Ap J, 487, 636
- Hartman, R. C., et al. 1992, in The Compton Observatory Science Workshop, ed. C. R. Shrader, N. Gehrels, & B. Dennis (NASA Conf. Publ 3137), 116
- Isobe, T., Feigelson, E. D., & Nelson, P. I. 1986, ApJ, 306, 490
- Isobe, T. & Feigelson, E. D. 1990, BAAS, 22, 917
- Jones, T. W., O’Dell, S. L., & Stein, W. A. 1974a, ApJ, 188, 353
- Jones, T. W., O’Dell, S. L., & Stein, W. A. 1974b, ApJ, 192, 261
- Kembhavi, A., Feigelson, E. D., & Singh, K. P. 1986, MNRAS, 220, 51
- Königl, A. 1981, ApJ, 243, 700
- Krichbaum, T. P., et al. 1990, A&A, 230, 271
- Ku, W. H.-M., Helfand, D. J., & Lucy, L. B. 1980, Nature, 288, 323
- Leach, C. M., Mc Hardy, I. M., & Papadakis, I. E. 1995, MNRAS, 272, 221
- Ledden, J. E., & O’Dell, S. L. 1985, ApJ, 298, 630

- Liszt, H. S., & Wilson, R. W. 1993, *ApJ*, 403, 663
- Malaguti, G., Bassani, L., & Caroli, E. 1994, *ApJS*, 94, 517
- Marscher, A. P. 1977, *ApJ*, 216, 244
- Marscher, A.P. 1980, *ApJ*, 235, 386
- Marscher, A. P. 1987 in *Superliminal Radio Sources*, ed. J. A. Zensus & T. J. Pearson (Cambridge Univ. Press) 280
- Marscher, A. P. 1988, *ApJ*, 334, 552
- Marscher, A. P., & Gear, W. K. 1985, *ApJ*, 298, 114
- Marscher, A. P., Gear, W. K., & Travis, J. P. 1992, in *Variability of Blazars*, ed. E. Valtaoja & M. Valtonen (Cambridge Univ. Press), 85
- Marscher, A. P., Zhang, Y. F., Shaffer, D. B., Aller, H. D., & Aller, M. F. 1991, *ApJ*, 371, 491
- Mattox, J. R., et al. 1993, *ApJ*, 410, 609
- Mattox, J. R., et al. 1996, *ApJ*, 461, 396
- Melia, F., & Königl, A. 1989, *ApJ*, 340, 162
- Owen, F. N., Helfand, D. J., & Spangler, S. R. 1981, *ApJ*, 250, L55
- Padovani, P. 1992, *A&A*, 256, 399
- Pearson, T. J. & Readhead, A. C. S. 1984, *ARA&A*, 22, 97
- Pearson, T. J. & Readhead, A. C. S. 1988, *ApJ*, 328, 114
- Petre, R., Mushotzsky, R. F., Krolik, J. H., & Holt, S. S. 1984, *ApJ*, 280, 499

- Sambruna, R. 1997, *Ap J*, 487, 536
- Schwab, F. R. & Cotton, W. D. 1983, *AJ*, 88, 688
- Stark, A. A., et al. 1992, *ApJS*, 79, 77
- Steppe, H., et al. 1988, *A&AS*, 75, 317
- Swanenburg, B. N., et al. 1978, *Nature*, 275, 298
- Tananbaum, H., et al. 1979, *ApJ*, 234, L9
- Thompson, D. J., et al. 1993, *ApJ*, 415, L13
- Thompson, D. J. , et al 1995. *ApJS*, 101, 259
- Thompson, D. J. , et al 1996, *ApJS*, 107, 227
- Tornikoski, M., et al. 1994, *A&A*, 289, 673
- Unwin, S. C., et al. 1994, *ApJ*, 432, 103
- Unwin, S. C., et al. 1997, *ApJ*, 480, 596
- von Linde, J., et al. 1993, *A&A*, 267, L23
- von Montigny, C., et al. 1995, *ApJ*, 440, 525
- Wehrle, A. E., & Mattox, J. R. 1994, in *The Second Compton Symposium*, ed. C. E. Fichtel, N. Gehrels, & J. P. Norris, *AIP Conf. Proc.* 304, 688
- Wilkes, B. J. 1992, *ROSAT-PROS Users Guide* (Cambridge, MA: Smithsonian Astrophys. Obs.)
- Wilkes, B. J., & Elvis, M. 1987, *ApJ*, 323, 243

Worrall, D. M., Giommi, P., Tananbaum, H., & Zamorani, G. 1987, ApJ, 313, 596

Worrall, D. M. and Wilkes, B. J. 1990, ApJ, 360, 396

Zamorani, G. et al. 1981, ApJ, 245, 357

Zhang, Y. F. 1994, PhD Dissertation, Boston University

Zhang, Y. F., & Marscher, A. P. 1994, in Proceedings of the 1993 ROSAT Science Symposium, ed. E. Schlegel & R. Petre (New York: AIP)

Zhou, Y. Y. et al 1997, ApJ, 484, L47

FIGURE CAPTIONS

Fig. 1.— VLBI images at 8.4 and 22 GHz from the Global VLBI Network. Parameters of each map are listed in Table 3. The restoring beam is shown in the lower left corner of each separate plot.

Fig. 2.— VLBA images at 22 GHz

Fig. 3.— Determination of the turnover frequency from VLBI and total flux density data. The dotted lines show the frequency ν_n at which the extrapolation of the partially opaque and optically thick spectra meet. The solid curve is the synchrotron spectrum for a homogeneous sphere with spectral turnover at frequency ν_m as discussed in the text.

Fig. 4.— The synchrotron spectrum (fit by a self-absorbed, uniform source model) and predicted first order (middle curve) and second order (rightmost curve) spectra for 0133+476, plotted against the contemporaneous data.

Fig. 5.— Similar to Figure 4, except for the quasar 0836+710. The EGRET gamma-ray data from Thompson et al. (1993) is plotted to the right on the 1992 March spectrum. Optical data are from von Linde et al. (1993). The large uncertainty in the optical point corresponds to the range of fluxes during a short term flare.

Fig. 6.— Correlation plot between VLBI core flux density at 22 GHz and X-ray flux density at 1 keV.

Fig. 7.— VLBI core luminosity at 22 GHz vs. X-ray luminosity at 1 keV. The solid line corresponds to a linear regression.

Fig. 8.— Millimeter-wave flux density (273 GHz) vs. X-ray flux density at 1 keV.

Fig. 9.— Millimeter-wave luminosity (273 GHz) vs. X-ray luminosity at 1 keV.

Fig. 10.— Flux density at the turnover frequency ν_m vs. X-ray flux density at 1 keV.

Fig. 11.— Luminosity at the turnover frequency ν_m vs. X-ray luminosity at 1 keV. The solid line is a linear regression.

Fig. 12.— X-ray spectral index vs. millimeter-wave spectral index.

Fig. 13.— Histogram of turnover frequencies ν_m .

Fig. 14.— Histogram of innermost radius measured by our 22GHz VLBI observations.

Fig. 15.— Histogram of X-ray to millimeter-wave spectral index, defined as $\alpha_{mmx} = \log(F_x/F_{\text{mm}})/\log(2.4 \times 10^{17}/2.7 \times 10^{11})$

Fig. 16.— X-ray to millimeter-wave spectral index α_{mmx} vs. radio core fraction at 8.4 GHz.

Fig. 17.— X-ray to millimeter-wave spectral index α_{mmx} vs. radio core fraction at 22 GHz.

TABLE 10
PHYSICAL PARAMETERS

Source (1)	$B(\frac{\delta_{min}}{\delta})$ (2)	$N_O(\frac{\delta}{\delta_{min}})^{2(\alpha+2)}$ (3)	R (4)	γ_{min} (5)	γ_{max} (6)	α_{1C} (7)
0016+731	41	1×10^{-11}	4.2	~ 10	3×10^3	1.37
0133+476	$\lesssim 6 \times 10^{-2}$	$\gtrsim 5 \times 10^{-6}$	$\lesssim 0.41$	~ 10	$\gtrsim 8 \times 10^3$	0.66
0212+735	17	6×10^{-12}	2.4	~ 10	5×10^3	1.59
NRAO 140	24	2.6×10^{-11}	3.1	~ 10	5×10^3	1.35
3C 111	21	5×10^{-5}	0.33	~ 10	4×10^3	0.76
3C 120	$\lesssim 20$	$\gtrsim 7 \times 10^{-5}$	$\lesssim 0.19$	~ 10	$\gtrsim 5 \times 10^3$	0.81
0552+398	5	2.3×10^{-9}	3.1	~ 10	4×10^3	1.06
0642+449	6	8.5×10^{-13}	4.6	~ 10	4×10^3	1.85
0736+017	$\lesssim 16$	$\gtrsim 7.2 \times 10^{-7}$	$\lesssim 0.75$	~ 10	$\gtrsim 3 \times 10^3$	0.93
0804+499	$\lesssim 2$	$\gtrsim 1 \times 10^{-6}$	$\lesssim 1.3$	~ 10	$\gtrsim 3 \times 10^3$	0.66
0836+710	$\lesssim 1 \times 10^{-1}$	$\gtrsim 1 \times 10^{-6}$	$\lesssim 1.1$	~ 100	$\gtrsim 1 \times 10^4$	0.76
0923+392	1×10^{-2}	2×10^{-5}	2.6	~ 10	2×10^4	0.54
0945+408	80	2.5×10^{-7}	3.5	~ 10	2×10^3	0.78
0955+476	47	6.3×10^{-8}	2.1	~ 10	3×10^3	0.99
1611+343	3	1×10^{-6}	1.6	~ 10	4×10^3	0.62
1633+382	140	6.5×10^{-6}	2.1	~ 10	2×10^3	0.68
3C 345	$\lesssim 1$	$\gtrsim 2 \times 10^{-6}$	$\lesssim 1.4$	~ 10	$\gtrsim 4 \times 10^3$	0.65
2136+141	$\lesssim 1 \times 10^{-1}$	$\gtrsim 3 \times 10^{-7}$	$\lesssim 2.8$	~ 10	$\gtrsim 1 \times 10^4$	0.78
2201+315	0.18	2.0×10^{-5}	1.2	~ 10	1×10^4	0.70

TABLE 11
CORRELATION AND REGRESSION PARAMETERS

Dependent (1)	Independent (2)	N (3)	% Censored (4)	Spearman (5)	Probability (6)	Slope (7)	Intercept (8)	Method (9)
$\log F_{\nu x}$	$\log F_{\nu c}$	21	0	-0.130	0.5614	...	...	...
$\log L_{\nu x}$	$\log L_{\nu c}$	21	0	0.789	0.0004	0.49 ± 0.09	10.91	BJ
$\log L_{\nu x}$	$\log L_{\nu c}$	19	0	0.717	0.0023	0.67 ± 0.14	4.35	BJ
$\log F_{\nu x}$	$\log F_{\nu mm}$	24	0	0.342	0.1007	0.38 ± 0.37	-0.38	BJ
$\log L_{\nu x}$	$\log L_{\nu mm}$	24	0	0.651	0.0018	0.60 ± 0.14	7.32	BJ
$\log L_{\nu x}$	$\log L_{\nu mm}$	22	0	0.548	0.0120	0.80 ± 0.24	0.72	BJ
$\log F_{\nu x}$	$\log F_{\nu m}$	24	17	0.342	0.1007	0.85 ± 0.45	-0.85	SB
$\log L_{\nu x}$	$\log L_{\nu m}$	24	17	0.670	0.0013	0.53 ± 0.10	5.27	SB
$\log L_{\nu x}$	$\log L_{\nu m}$	22	18	0.570	0.0091	0.73 ± 0.17	2.56	SB
$\log F_{\nu x}$	$\log \theta$	21	29	-0.234	0.2954	...	...	...
$\log F_{\nu x}$	$\log \nu_m$	24	19	0.168	0.4300	...	...	...
$\log L_{\nu x}$	z	24	0	0.741	0.0004	...	...	...
$\log L_{\nu c}$	z	21	0	0.917	< 0.0001	...	...	...
$\log L_{\nu m}$	z	24	17	0.857	< 0.0001	...	...	...
$\log L_{\nu mm}$	z	24	0	0.705	0.0007	...	...	...
α_x	α_{mm}	24	0	-0.163	0.4343	...	...	...
α_x	α_{ir}	13	0	-0.335	0.2463	...	...	...
α_{ir}	α_{mm}	15	0	-0.125	0.6394	...	...	...
θ	$\log \nu_m$	24	42	0.127	0.5146	...	...	...